

**BCKDK Antibody**  
**Rabbit mAb**  
**Catalog # AP92905****Specification****BCKDK Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O14874</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
BCKD kinase; BCKDHKIN; Bckdk; BCKDKD; BDK;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	46360 Da

**BCKDK Antibody - Additional Information**

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human BCKDK
Description	Catalyzes the phosphorylation and inactivation of the branched-chain alpha-ketoacid dehydrogenase complex, the key regulatory enzyme of the valine, leucine and isoleucine catabolic pathways. Key enzyme that regulate the activity state of the BCKD complex.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**BCKDK Antibody - Protein Information****Name** BCKDK {ECO:0000303|PubMed:29779826, ECO:0000312|HGNC:HGNC:16902}**Function**

Serine/threonine-protein kinase component of macronutrients metabolism. Forms a functional kinase and phosphatase pair with PPM1K, serving as a metabolic regulatory node that coordinates branched-chain amino acids (BCAAs) with glucose and lipid metabolism via two distinct phosphoprotein targets: mitochondrial BCKDHA subunit of the branched- chain alpha-ketoacid dehydrogenase (BCKDH) complex and cytosolic ACLY, a lipogenic enzyme of Krebs cycle (PubMed:<a href="http://www.uniprot.org/citations/24449431" target="\_blank">24449431</a>, PubMed:<a href="http://www.uniprot.org/citations/29779826" target="\_blank">29779826</a>, PubMed:<a href="http://www.uniprot.org/citations/37558654" target="\_blank">37558654</a>).

Phosphorylates and inactivates mitochondrial BCKDH complex a multisubunit complex consisting of three multimeric components each involved in different steps of BCAA catabolism: E1 composed of BCKDHA and BCKDHB, E2 core composed of DBT monomers, and E3 composed of DLD monomers. Associates with the E2 component of BCKDH complex and phosphorylates BCKDHA on Ser-337, leading to conformational changes that interrupt substrate channeling between E1 and E2 and inactivates the BCKDH complex (PubMed:<a href="http://www.uniprot.org/citations/29779826" target="\_blank">29779826</a>, PubMed:<a href="http://www.uniprot.org/citations/37558654" target="\_blank">37558654</a>). Phosphorylates ACLY on Ser-455 in response to changes in cellular carbohydrate abundance such as occurs during fasting to feeding metabolic transition. Refeeding stimulates MLXIPL/ChREBP transcription factor, leading to increased BCKDK to PPM1K expression ratio, phosphorylation and activation of ACLY that ultimately results in the generation of malonyl-CoA and oxaloacetate immediate substrates of de novo lipogenesis and glucogenesis, respectively (PubMed:<a href="http://www.uniprot.org/citations/29779826" target="\_blank">29779826</a>). Recognizes phosphosites having SxxE/D canonical motif (PubMed:<a href="http://www.uniprot.org/citations/29779826" target="\_blank">29779826</a>).

#### Cellular Location

Mitochondrion matrix {ECO:0000250|UniProtKB:Q00972, ECO:0000305|PubMed:24449431}  
Note=Detected in the cytosolic compartment of liver cells {ECO:0000250|UniProtKB:Q00972}

#### Tissue Location


Ubiquitous.


#### BCKDK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### BCKDK Antibody - Images





Western blot analysis of BCKDK expression in HeLa cell lysate.