

OXGR1 / GPR99 Antibody
Rabbit mAb
Catalog # AP92922**Specification****OXGR1 / GPR99 Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	Q96P68
Reactivity	Rat
Clonality	Monoclonal
Other Names	
GPR80; GPR99; Oxgr1; P2RY15; P2Y15;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	38251 Da

OXGR1 / GPR99 Antibody - Additional Information

Dilution	WB~~1:1000 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human OXGR1 / GPR99
Description	Receptor for alpha-ketoglutarate. Seems to act exclusively through a G(q)-mediated pathway.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

OXGR1 / GPR99 Antibody - Protein Information**Name** OXGR1**Function**

G protein-coupled receptor for dicarboxylates and amino dicarboxylates (PubMed: 15141213, PubMed: 36571463, PubMed: 36919698). Receptor for itaconate, a metabolite produced by myeloid lineages (PubMed: 36919698). In the respiratory epithelium, couples the binding of itaconate to the activation of GNA11 and downstream intracellular Ca(2+) release, leading to mucocilliary clearance of airborne pathogens (PubMed: 36919698). Receptor for leukotriene E4 (LTE4) produced by mast cells upon allergic inflammation. Binds with

high affinity to LTE4 and elicits mucin release from pulmonary epithelium in response to airborne fungi allergens. Regulates mucin-producing goblet cell homeostasis (By similarity). Receptor for alpha-ketoglutarate produced by proximal tubule renal cells upon metabolic alkalosis. In an intrarenal paracrine signaling pathway, binds alpha-ketoglutarate and drives transepithelial salt reabsorption and bicarbonate secretion by SLC26A4/pendrin-positive intercalated cells (By similarity) (PubMed:15141213).

Cellular Location

Cell membrane; Multi-pass membrane protein. Note=Upon itaconate binding, internalizes via endocytosis in a beta-arrestin dependent manner

Tissue Location


Detected in kidney and, to a lower extent, in placenta. Not detected in brain tissues including the frontal cortex, caudate putamen, thalamus, hypothalamus, hippocampus or pons

OXGR1 / GPR99 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OXGR1 / GPR99 Antibody - Images



Western blot analysis of OXGR1 / GPR99 expression in A549 cell lysate.