

UCL5IP Antibody
Rabbit mAb
Catalog # AP93042**Specification**

UCL5IP Antibody - Product Information

Application	WB
Primary Accession	Q99871
Clonality	Monoclonal
Other Names	
Haus7; HSXQ28ORF; UCH37 interacting protein 1; UCL5IP; UIP1;	

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	40778 Da

UCL5IP Antibody - Additional Information

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human UCL5IP
Description	Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

UCL5IP Antibody - Protein Information**Name** HAUS7**Synonyms** UCL5IP, UIP1**Function**

Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Localizes to interphase centrosomes and to mitotic spindle microtubules.

Tissue Location


Detected in spleen, thymus, testis, ovary, small intestine and colon, with highest levels of expression in testis and ovary.

UCHL5IP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UCHL5IP Antibody - Images



Western blot analysis of UCHL5IP expression in HeLa cell lysate.