

CRCP Antibody
Rabbit mAb
Catalog # AP93061**Specification**

CRCP Antibody - Product Information

Application	WB, IHC, ICC, IP
Primary Accession	075575
Reactivity	Rat
Clonality	Monoclonal
Other Names	
CGRPRCP; CRCP; HsC17; RCP; RCP9;	

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	16871 Da

CRCP Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CRCP
Description	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Specific peripheric component of RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage Condition and Buffer	

CRCP Antibody - Protein Information**Name** CRCP ([HGNC:17888](#))**Function**

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates (PubMed:20413673, PubMed:33558764, PubMed:34675218). Specific peripheric component of RNA polymerase III (Pol III) which synthesizes small non-coding RNAs

including 5S rRNA, snRNAs, tRNAs and miRNAs from at least 500 distinct genomic loci. With POLR3H/RPC8 forms a mobile stalk that protrudes from Pol III core and functions primarily in transcription initiation (By similarity) (PubMed:20413673, PubMed:33558764, PubMed:33558766, PubMed:34675218). Pol III plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF-kappa-B through the RIG-I pathway (PubMed:19609254, PubMed:19631370).

Cellular Location

Nucleus. Cell membrane {ECO:0000250|UniProtKB:O35427}; Peripheral membrane protein {ECO:0000250|UniProtKB:O35427}; Cytoplasmic side {ECO:0000250|UniProtKB:O35427}

Tissue Location


Ubiquitous. Most prevalent in testis.

CRCP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRCP Antibody - Images



Western blot analysis of CRCP expression in HeLa cell lysate.