

IMP4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9358b

Specification

IMP4 Antibody (C-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q96G21
Other Accession	Q5POR5 , Q8VHZ7
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33757
Antigen Region	215-243

IMP4 Antibody (C-term) - Additional Information

Gene ID 92856

Other Names

U3 small nucleolar ribonucleoprotein protein IMP4, U3 snoRNP protein IMP4, Brix domain-containing protein 4, IMP4, BXDC4

Target/Specificity

This IMP4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 215-243 amino acids from the C-terminal region of human IMP4.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IMP4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IMP4 Antibody (C-term) - Protein Information

Name IMP4 ([HGNC:30856](#))

Synonyms BXDC4

Function Component of the 60-80S U3 small nucleolar ribonucleoprotein (U3 snoRNP). Required for the early cleavages during pre-18S ribosomal RNA processing (PubMed:[12655004](#)). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:[34516797](#)).

Cellular Location


Nucleus, nucleolus

IMP4 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IMP4 Antibody (C-term) - Images



All lanes : Anti-IMP4 Antibody (C-term) at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



IMP4 Antibody (C-term) (Cat. #AP9358b) IHC analysis in formalin fixed and paraffin embedded human testis carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the IMP4 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

IMP4 Antibody (C-term) - Background

IMP4 forms a ternary complex with IMP3 (MIM 612980) and MPP10 (MPHOSPHO10; MIM 605503) that interacts with U3 small nucleolar RNA (snoRNA), which is required for the early cleavage steps in pre-rRNA processing.

IMP4 Antibody (C-term) - References

Hillier, L.W. Nature 434 (7034), 724-731 (2005)
Andersen, J.S. Nature 433 (7021), 77-83 (2005)
Granneman, S. Nucleic Acids Res. 31 (7), 1877-1887 (2003)