

## Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody

**Catalog # AP93610** 

#### **Specification**

# Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Clonality WB P68431/Q71DI3/P84243 Human Polyclonal

### Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Additional Information

**Storage Conditions** -20°C

Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Protein Information

## Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cvtometv
- Cell Culture

## Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Images



Western blot analysis of Hela, diluted at 1:1000. Secondary antibody was diluted at 1:20000 cells





Tel: 858.875.1900 Fax: 858.875.1999

nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit.

## Histone H3 (Di Methyl Lys18) Rabbit Polyclonal Antibody - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015],