

LIPE Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9376b**Specification**

LIPE Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q05469
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	116598
Antigen Region	822-849

LIPE Antibody (C-term) - Additional Information**Gene ID** 3991**Other Names**

Hormone-sensitive lipase, HSL, LIPE

Target/Specificity

This LIPE antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 822-849 amino acids from the C-terminal region of human LIPE.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LIPE Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

LIPE Antibody (C-term) - Protein Information**Name** LIPE

Function Lipase with broad substrate specificity, catalyzing the hydrolysis of triacylglycerols (TAGs), diacylglycerols (DAGs), monoacylglycerols (MAGs), cholesteryl esters and retinyl esters (PubMed:[15716583](#), PubMed:[15955102](#), PubMed:[19800417](#), PubMed:[8812477](#)). Shows a preferential hydrolysis of DAGs over TAGs and MAGs and preferentially hydrolyzes the fatty acid (FA) esters at the sn-3 position of the glycerol backbone in DAGs (PubMed:[19800417](#)). Preferentially hydrolyzes FA esters at the sn-1 and sn-2 positions of the glycerol backbone in TAGs (By similarity). Catalyzes the hydrolysis of 2-arachidonoylglycerol, an endocannabinoid and of 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor (By similarity). In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production (By similarity).

Cellular Location

Cell membrane. Membrane, caveola. Cytoplasm, cytosol. Lipid droplet {ECO:0000250|UniProtKB:P54310}. Note=Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation (PubMed:17026959). Phosphorylation by AMPK reduces its translocation towards the lipid droplets (By similarity) {ECO:0000250|UniProtKB:P54310, ECO:0000269|PubMed:17026959}

Tissue Location


Testis..

LIPE Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


LIPE Antibody (C-term) - Images



Western blot analysis of LIPE Antibody (C-term) (Cat. #AP9376b) in HL-60 cell line lysates (35ug/lane). LIPE (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human testis tissue reacted with LIPE Antibody (C-term) (Cat. #AP9376b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



LIPE Antibody (C-term) (Cat. #AP9376b) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

LIPE Antibody (C-term) - Background

LIPE has a long and a short form, generated by use of alternative translational start codons. The long form is expressed in steroidogenic tissues such as testis, where it converts cholesteryl esters to free cholesterol for steroid hormone production. The short form is expressed in adipose tissue, among others, where it hydrolyzes stored triglycerides to free fatty acids.

LIPE Antibody (C-term) - References

- Bezaire, V., et al. FEBS Lett. 583(18):3045-3049(2009)
- Chen, H.H., et al. Am. J. Clin. Nutr. 90(2):255-262(2009)
- Kuzmin, A., et al. Biol. Reprod. 81(2):319-326(2009)
- Bezaire, V., et al. J. Biol. Chem. 284(27):18282-18291(2009)
- Drenos, F., et al. Hum. Mol. Genet. 18(12):2305-2316(2009)