

FXYD3 (15L6) Mouse Monoclonal antibody
FXYD3 (15L6) Mouse Monoclonal antibody
Catalog # AP93862**Specification**

FXYD3 (15L6) Mouse Monoclonal antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC, IF |
| Primary Accession | Q14802 |
| Reactivity | Human |
| Clonality | Monoclonal |
| Calculated MW | 9263 |

FXYD3 (15L6) Mouse Monoclonal antibody - Additional Information**Gene ID** 5349**Other Names**

FXYD domain-containing ion transport regulator 3, Chloride conductance inducer protein Mat-8, Mammary tumor 8 kDa protein, Phospholemman-like, Sodium/potassium-transporting ATPase subunit FXYD3, FXYD3, MAT8, PLML

Dilution

WB~~1:1000
IHC~~1:100~500
IF~~1:50~200

Storage Conditions

-20°C

FXYD3 (15L6) Mouse Monoclonal antibody - Protein Information**Name** FXYD3**Synonyms** MAT8, PLML**Function**

Associates with and regulates the activity of the sodium/potassium-transporting ATPase (NKA) which transports Na(+) out of the cell and K(+) into the cell (PubMed:17077088). Reduces glutathionylation of the NKA beta-1 subunit ATP1B1, thus reversing glutathionylation-mediated inhibition of ATP1B1 (PubMed:21454534). Induces a hyperpolarization-activated chloride current when expressed in Xenopus oocytes (PubMed:7836447).

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Isoform 1: Expressed mainly in differentiated cells (at protein level). Isoform 2: Expressed mainly in undifferentiated cells (at protein level).


FXYD3 (15L6) Mouse Monoclonal antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FXYD3 (15L6) Mouse Monoclonal antibody - Images


Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-FXYD3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93862)




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-FXYD3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93862)




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-FXYD3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-FXYD3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93862)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-FXYD3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93862)



Anti-FXYD3 mouse monoclonal antibody (AP93862) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY FXYD3 .



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-FXYD3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93862)

FXYD3 (15L6) Mouse Monoclonal antibody - Background

This gene belongs to a small family of FXYD-domain containing regulators of Na⁺/K⁺ ATPases which share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD, and containing 7 invariant and 6 highly conserved amino acids. This gene encodes a cell membrane protein that may regulate the function of ion-pumps and ion-channels. This gene may also play a role in tumor progression. Alternative splicing results in multiple transcript variants encoding distinct isoforms.