

phospho-FADD (Ser194) Rabbit pAb
phospho-FADD (Ser194) Rabbit pAb
Catalog # AP93921**Specification**

phospho-FADD (Ser194) Rabbit pAb - Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23 KDa
Physical State	Liquid
Immunogen	KLH conjugated Synthesised phosphopeptide derived from human FADD around the phosphorylation site of Ser194
Epitope Specificity	AM(p-S)PM
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 death domain. Contains 1 DED (death effector) domain.
SUBUNIT	Can self-associate. Interacts with CFLAR, PEA15 and MBD4. When phosphorylated, part of a complex containing HIPK3 and FAS. May interact with MAVS/IPS1. Interacts with MOCV v-CFLAR protein and LRDD. Interacts (via death domain) with FAS (via death domain). Interacts with CASP8.
DISEASE	The interaction between the FAS and FADD death domains is crucial for the formation of the death-inducing signaling complex (DISC). Defects in FADD are the cause of infections recurrent associated with encephalopathy hepatic dysfunction and cardiovascular malformations (IEHDCM) [MIM:613759]. A condition with biological features of autoimmune lymphoproliferative syndrome such as high-circulating CD4(-)CD8(-)TCR-alpha-beta(+) T-cell counts, and elevated IL10 and FASL levels. Affected individuals suffer from recurrent, stereotypical episodes of fever, encephalopathy, and mild liver dysfunction sometimes accompanied by generalized seizures. The episodes can be triggered by varicella zoster virus (VZV), measles

Important Note

mumps rubella (MMR) attenuated vaccine, parainfluenza virus, and Epstein-Barr virus (EBV).

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Predicted to enable several functions, including caspase binding activity; death effector domain binding activity; and tumor necrosis factor receptor superfamily binding activity. Involved in several processes, including hematopoietic or lymphoid organ development; negative regulation of activation-induced cell death of T cells; and positive regulation of CD8-positive, alpha-beta cytotoxic T cell extravasation. Acts upstream of or within extrinsic apoptotic signaling pathway in absence of ligand; motor neuron apoptotic process; and regulation of programmed cell death. Predicted to be located in several cellular components, including cell body; cytosol; and membrane raft. Predicted to be part of CD95 death-inducing signaling complex and ripoptosome. Predicted to be active in cytoplasm. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Human ortholog(s) of this gene implicated in leukemia. Orthologous to human FADD (Fas associated via death domain). [provided by Alliance of Genome Resources, Apr 2022]

phospho-FADD (Ser194) Rabbit pAb - Additional Information**Target/Specificity**

Expressed in a wide variety of tissues, except for peripheral blood mononuclear leukocytes.

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

phospho-FADD (Ser194) Rabbit pAb - Protein Information**phospho-FADD (Ser194) Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

phospho-FADD (Ser194) Rabbit pAb - Images



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-FADD (Ser194)) Polyclonal Antibody, Unconjugated (AP93921) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-FADD (Ser194)) Polyclonal Antibody, Unconjugated (AP93921) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

phospho-FADD (Ser194) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.