

**IRS3 Rabbit pAb**  
**IRS3 Rabbit pAb**  
**Catalog # AP93940****Specification**

---

**IRS3 Rabbit pAb - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | <b>IHC-P, WB</b>       |
| Primary Accession | <a href="#">O54718</a> |
| Reactivity        | <b>Mouse</b>           |
| Host              | <b>Rabbit</b>          |
| Clonality         | <b>Polyclonal</b>      |

**IRS3 Rabbit pAb - Additional Information****Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


**Storage**


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**IRS3 Rabbit pAb - Protein Information****IRS3 Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**IRS3 Rabbit pAb - Images**



Tissue/cell: human pneumonia tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-IRS-3 Polyclonal Antibody, Unconjugated(AP93940) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample: Lane 1: Adipose (Rat) Lysate at 40 ug Primary: Anti-IRS3 (AP93940) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kD  
Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide

for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRS3) Polyclonal Antibody, Unconjugated (AP93940) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**IRS3 Rabbit pAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.