

phospho-HSP70 (Tyr41) Rabbit pAb phospho-HSP70 (Tyr41) Rabbit pAb Catalog # AP93958

# Specification

## phospho-HSP70 (Tyr41) Rabbit pAb - Product Information

Application
Primary Accession

Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

**Epitope Specificity** 

Isotype **Purity** 

affinity purified by Protein A

Polycional

Human Rabbit Polyclonal 70 KDa Liquid

PODMV9

**KLH conjugated Synthesised** 

WB, IHC-P, IHC-F, IF

phosphopeptide derived from human HSP70 around the phosphorylation site of

Tyr41 PS(p-Y)VA

IgG

**Buffer** 

SUBCELLULAR LOCATION

**SIMILARITY** 

**SUBUNIT** 

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated

mRNAs.

Belongs to the heat shock protein 70

family.

Component of the CatSper complex. Identified in a mRNP granule complex, at least composed of ACTB, ACTN4, DHX9, ERG, HNRNPA1, HNRNPA2B1, HNRNPAB, HNRNPD, HNRNPL, HNRNPR, HNRNPU, HSPA1, HSPA8, IGF2BP1, ILF2, ILF3, NCBP1, NCL, PABPC1, PABPC4, PABPN1, RPLPO. RPS3. RPS3A. RPS4X. RPS8. RPS9. SYNCRIP, TROVE2, YBX1 and untranslated mRNAs. Interacts with TSC2. Interacts with IRAK1BP1. Interacts with TERT: the interaction occurs in the absence of the RNA component, TERC, and dissociates once the TERT complex has formed. Interacts with DNAJC7. Interacts with CHCHD3.

Important Note

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

#### **Background Descriptions**

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes existing



proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins. [provided by RefSeq, Jul 2008].

## phospho-HSP70 (Tyr41) Rabbit pAb - Additional Information

Gene ID 3303;3304

#### **Other Names**

Heat shock 70 kDa protein 1B {ECO:0000312|HGNC:HGNC:5233}, Heat shock 70 kDa protein 2, HSP70-2, HSP70.2, Heat shock protein family A member 1B, HSPA1B (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=5233" target="blank">HGNC:5233</a>)

### Target/Specificity

HSPA1B is testis-specific.

#### **Dilution**

```
<span class ="dilution_WB">WB~~1:1000</span><br \> <span class
="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class = "dilution_IF">IF~~1:50~200</span>
```

## Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### phospho-HSP70 (Tyr41) Rabbit pAb - Protein Information

Name HSPA1B (HGNC:5233)

## **Function**

Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The co- chaperones have been shown to not only regulate different steps of the ATPase cycle, but they also have an individual specificity such that one co-chaperone may promote folding of a substrate while another may promote degradation. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release. The co-chaperones are of three types: J-domain co-chaperones such as HSP40s (stimulate ATPase hydrolysis by HSP70), the nucleotide exchange factors (NEF) such as BAG1/2/3 (facilitate conversion of HSP70 from the ADP-bound to the ATP-bound state thereby promoting substrate release), and the TPR domain chaperones such as HOPX and STUB1 (PubMed: <a  $href="http://www.uniprot.org/citations/24012426" \ target="\_blank">24012426</a>, PubMed:<a href="http://www.uniprot.org/citations/24318877" target="_blank">24012426</a>, PubMed:<a href="http://www.uniprot.org/citations/24318877" target="_blank">24318877</a>, PubMed:<a href="http://www.uniprot.org/citations/2431887" target="_blank">14318878</a>, PubMed$ href="http://www.uniprot.org/citations/26865365" target="\_blank">26865365</a>). Maintains



protein homeostasis during cellular stress through two opposing mechanisms: protein refolding and degradation. Its acetylation/deacetylation state determines whether it functions in protein refolding or protein degradation by controlling the competitive binding of co-chaperones HOPX and STUB1. During the early stress response, the acetylated form binds to HOPX which assists in chaperone-mediated protein refolding, thereafter, it is deacetylated and binds to ubiquitin ligase STUB1 that promotes ubiquitin-mediated protein degradation (PubMed:<a href="http://www.uniprot.org/citations/27708256" target="\_blank">27708256</a>). Regulates centrosome integrity during mitosis, and is required for the maintenance of a functional mitotic centrosome that supports the assembly of a bipolar mitotic spindle (PubMed:<a href="http://www.uniprot.org/citations/27137183" target="\_blank">27137183</a>(a>). Enhances STUB1-mediated SMAD3 ubiquitination and degradation and facilitates STUB1-mediated inhibition of TGF-beta signaling (PubMed:<a href="http://www.uniprot.org/citations/24613385" target="\_blank">24613385</a>(a>). Essential for STUB1-mediated ubiquitination and degradation of FOXP3 in regulatory T-cells (Treg) during inflammation (PubMed:<a href="http://www.uniprot.org/citations/23973223" target=" blank">23973223</a>(a>).

#### **Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs

### **Tissue Location**


HSPA1B is testis-specific.

#### phospho-HSP70 (Tyr41) Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# phospho-HSP70 (Tyr41) Rabbit pAb - Images






Blank control (blue line): Jurkat (fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody (green line): Rabbit Anti-phospho-HSP70 (Tyr41) antibody (AP93958),Dilution: 1  $\mu$ g /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC,Dilution: 1  $\mu$ g /test.




Tissue/cell: Rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-phospho-HSP70(Tyr41)Polyclonal Antibody, Unconjugated(AP93958) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Blank control: MCF7. Primary Antibody (green line): Rabbit Anti- phospho-HSP70 (Tyr41) antibody (AP93958) Dilution: 2  $\mu$ g /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 1  $\mu$ g /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.






Blank control:A549. Primary Antibody (green line): Rabbit Anti-phospho-HSP70 (Tyr41) antibody (AP93958) Dilution: 1  $\mu$ g /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-AF488 Dilution: 1  $\mu$ g /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Sample: Epithelium (Mouse) Lysate at 40 ug Primary: Anti-phospho-HSP70 (Tyr41) (AP93958) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 70 kD





Sample: A549(Human) Cell Lysate at 30 ug K562(Human) Cell Lysate at 30 ug Primary: Anti-phospho-HSP70 (Tyr41) (AP93958) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 70 kD

# phospho-HSP70 (Tyr41) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.