

Neuronal thread protein AD7c-NTP Rabbit pAb

Neuronal thread protein AD7c-NTP Rabbit pAb Catalog # AP93969

Specification

Neuronal thread protein AD7c-NTP Rabbit pAb - Product Information

Application WB, IHC-P, IHC-F, IF

Reactivity
Host
Clonality
Calculated MW
Physical State
Human
Rabbit
Polyclonal
41 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human Neuronal thread protein

AD7c-NTP 301-375/375

IgG

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

AD7c-NTP is detected in increased concentration in the cortical neurons, brain-tissue extracts, cerebrospinal fluid, and urine early in the course of AD neurodegeneration, and its level is positively correlated with the severity of dementia. All these characteristics make it a possible biomarker for AD.

Neuronal thread protein AD7c-NTP Rabbit pAb - Additional Information

Dilution

WB~~1:1000<br \><span class</pre>

="dilution IHC-P">IHC-P \sim N/A
<span class

="dilution_IHC-F">IHC-F~~N/A
span class ="dilution_IF">IF~~1:50~200

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Neuronal thread protein AD7c-NTP Rabbit pAb - Protein Information


Neuronal thread protein AD7c-NTP Rabbit pAb - Protocols




Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Neuronal thread protein AD7c-NTP Rabbit pAb - Images




Tissue/cell: Mouse brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-AD7c-NTP Polyclonal Antibody, Unconjugated(AP93969) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Tissue/cell: human brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-AD7c-NTP Polyclonal Antibody, Unconjugated(AP93969) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining






Sample: Brain (Mouse) Lysate at 40 ug Primary: Anti- AD7c-NTP(AP93969) at 1/300 dilution Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution Predicted band size: 41 kD Observed band size: 41 kD




Blank control: SHSY5Y. Primary Antibody (green line): Rabbit Anti-Neuronal thread protein AD7c-NTP antibody (AP93969) Dilution: 1ug/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.





Paraformaldehyde-fixed, paraffin embedded (rat cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Neuronal thread protein AD7c-NTP) Polyclonal Antibody, Unconjugated (AP93969) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Rat Cerebellum tissue lysates Primary: Anti-Neuronal thread protein AD7c-NTP (AP93969) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kDa Observed band size: 43 kDa

Neuronal thread protein AD7c-NTP Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.