

HPV16 E6 + HPV18 E6 Rabbit pAb
HPV16 E6 + HPV18 E6 Rabbit pAb
Catalog # AP93982**Specification**

HPV16 E6 + HPV18 E6 Rabbit pAb - Product Information

Application	IHC-P, WB
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

HPV16 E6 + HPV18 E6 Rabbit pAb - Additional Information**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HPV16 E6 + HPV18 E6 Rabbit pAb - Protein Information**HPV16 E6 + HPV18 E6 Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HPV16 E6 + HPV18 E6 Rabbit pAb - Images



Tissue/cell: human colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-HPV16 E6+HPV18 E6 Polyclonal Antibody, Unconjugated(AP93982) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample: Lane 1: Recombinant HPV16 E6 protein, DsbC & His(bs-49102P) 0.1ug Lane 2: Recombinant HPV16 E6 protein, DsbC & His(bs-49102P) 0.05ug Primary: Anti-HPV16 E6 + HPV18 E6 (AP93982) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 11 kDa Observed band size: 50 kDa



Blank control□black line□:Hela. Primary Antibody (green line): Rabbit Anti-HPV16 E6 + HPV18 E6

antibody (AP93982) Dilution:1ug/Test; Secondary Antibody□white blue line□: Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control□orange line□: Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

HPV16 E6 + HPV18 E6 Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.