

**MUC6(ready to use) Mouse mAb**  
**MUC6(ready to use) Mouse mAb**  
**Catalog # AP94075****Specification**

---

**MUC6(ready to use) Mouse mAb - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">Q6W4X9</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	257051

**MUC6(ready to use) Mouse mAb - Additional Information****Gene ID** 4588**Other Names**

Mucin-6, MUC-6, Gastric mucin-6, MUC6

**Dilution**

IHC-P~~N/A  
IHC-F~~N/A  
IF~~1:50~200  
ICC~~N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**MUC6(ready to use) Mouse mAb - Protein Information****Name** MUC6**Function**

May provide a mechanism for modulation of the composition of the protective mucus layer related to acid secretion or the presence of bacteria and noxious agents in the lumen. Plays an important role in the cytoprotection of epithelial surfaces and are used as tumor markers in a variety of cancers. May play a role in epithelial organogenesis.

**Cellular Location**

Secreted.

**Tissue Location**

Expressed in the regenerative zone of gastric antrum, gastric body mucosa and gastric incisura mucosa. Expressed in the deeper mucous glands of gastric antrum. Overexpressed in Helicobacter


pylori infected gastric epithelium. Highly expressed in duodenal Brunner's glands, gall bladder, seminal vesicle, pancreatic centroacinar cells and ducts, and periductal glands of the common bile duct.

### **MUC6(ready to use) Mouse mAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **MUC6(ready to use) Mouse mAb - Images**




Paraformaldehyde-fixed, paraffin embedded (human gastric); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUC6) Monoclonal Antibody, Unconjugated (AP94075) overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUC6) Monoclonal Antibody, Unconjugated (AP94075) overnight at 4°C, followed

by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human gastric); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUC6) Monoclonal Antibody, Unconjugated (AP94075) overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.

#### **MUC6(ready to use) Mouse mAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.