

HBA2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9414C**Specification**

HBA2 Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	P69905
Other Accession	P21767
Reactivity	Human, Mouse
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	100-128

HBA2 Antibody (Center) - Additional Information**Gene ID** 3039;3040**Other Names**

Hemoglobin subunit alpha, Alpha-globin, Hemoglobin alpha chain, HBA1

Target/Specificity

This HBA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 100-128 amino acids from the Central region of human HBA2.

Dilution

WB~~1:2000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HBA2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HBA2 Antibody (Center) - Protein Information**Name** HBA1

Function Involved in oxygen transport from the lung to the various peripheral tissues.

Tissue Location


Red blood cells.

HBA2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HBA2 Antibody (Center) - Images



Western blot analysis of HBA2 Antibody (Center) (Cat. #AP9414c) in mouse liver tissue lysates (35ug/lane). HBA2 (arrow) was detected using the purified Pab.



Anti-HBA2 Antibody (Center) at 1:2000 dilution + human fetal heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 15 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing K562 cells stained with AP9414c (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP9414c, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

HBA2 Antibody (Center) - Background

HBA2 located on chromosome 16 spans about 30 kb and includes seven loci: 5'- zeta - pseudozeta - mu - pseudoalpha-1 - alpha-2 - alpha-1 - theta - 3'. The alpha-2 (HBA2) and alpha-1 (HBA1) coding sequences are identical. These genes differ slightly over the 5' untranslated regions and the introns, but they differ significantly over the 3' untranslated regions. Two alpha chains plus two beta chains constitute HbA, which in normal adult life comprises about 97% of the total hemoglobin; alpha chains combine with delta chains to constitute HbA-2, which with HbF (fetal hemoglobin) makes up the remaining 3% of adult hemoglobin. Alpha thalassemias result from deletions of each of the alpha genes as well as deletions of both HBA2 and HBA1; some nondeletion alpha thalassemias have also been reported.

HBA2 Antibody (Center) - References

Sessa, R., et al. Am. J. Hematol. 85(2):143-144(2010)
Sharma, V., et al. Hematology 14(5):297-300(2009)
Waye, J.S., et al. Hemoglobin 33(6):519-522(2009)
Roy, P., et al. Hemoglobin 33(6):486-491(2009)
Joly, P., et al. Hemoglobin 33(3):196-205(2009)