

#### **HSP70 Mouse mAb**

**HSP70** Mouse mAb **Catalog # AP94160** 

### **Specification**

#### **HSP70 Mouse mAb - Product Information**

Application **Primary Accession** Reactivity

Host Clonality Calculated MW Physical State

Immunogen

Isotype **Purity** 

affinity purified by Protein G

Buffer

SUBCELLULAR LOCATION

**SIMILARITY** 

**SUBUNIT** 

Important Note

WB, IHC-P, IHC-F, IF

PODMV9 Human Rabbit **Monoclonal 70 KDa** Liquid

KLH conjugated synthetic peptide derived

from human Hsp70

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated

mRNAs.

Belongs to the heat shock protein 70

family.

Component of the CatSper complex. Identified in a mRNP granule complex, at least composed of ACTB, ACTN4, DHX9, ERG, HNRNPA1, HNRNPA2B1, HNRNPAB, HNRNPD, HNRNPL, HNRNPR, HNRNPU, HSPA1, HSPA8, IGF2BP1, ILF2, ILF3, NCBP1, NCL, PABPC1, PABPC4, PABPN1, RPLPO, RPS3, RPS3A, RPS4X, RPS8, RPS9, SYNCRIP, TROVE2, YBX1 and untranslated mRNAs. Interacts with TSC2. Interacts with IRAK1BP1. Interacts with TERT: the interaction occurs in the absence of the RNA component, TERC, and dissociates once the TERT complex has formed. Interacts with DNAJC7. Interacts with CHCHD3.

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

# **Background Descriptions**

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility



complex class III region, in a cluster with two closely related genes which encode similar proteins. [provided by RefSeq, Jul 2008].

#### **HSP70 Mouse mAb - Additional Information**

Gene ID 3303;3304

### **Other Names**

Heat shock 70 kDa protein 1B {ECO:0000312|HGNC:HGNC:5233}, Heat shock 70 kDa protein 2, HSP70-2, HSP70.2, Heat shock protein family A member 1B, HSPA1B (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=5233" target="blank">HGNC:5233</a>)

#### Target/Specificity

HSPA1B is testis-specific.

#### **Dilution**

```
 < span class = "dilution_WB">WB~~1:1000 < /span> < br \> < span class = "dilution_IHC-P">IHC-P~~N/A < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~N/A < /span> < br \> < span class = "dilution_IF">IF~~1:50~200 < /span> < br \> < span class = "dilution_IF">IF~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~N/A < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IF">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < /span> < span class = "dilution_IHC-F">IHC-F~~1:50~200 < span class = "dil
```

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **HSP70 Mouse mAb - Protein Information**

Name HSPA1B (HGNC:5233)

# **Function**

Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The co-chaperones have been shown to not only regulate different steps of the ATPase cycle, but they also have an individual specificity such that one co-chaperone may promote folding of a substrate while another may promote degradation. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release. The co-chaperones are of three types: J-domain co-chaperones such as HSP40s (stimulate ATPase hydrolysis by HSP70), the nucleotide exchange factors (NEF) such as BAG1/2/3 (facilitate conversion of HSP70 from the ADP-bound to the ATP-bound state thereby promoting substrate release), and the TPR domain chaperones such as HOPX and STUB1 (PubMed: <a href="http://www.uniprot.org/citations/24012426" target=" blank">24012426</a>, PubMed:<a href="http://www.uniprot.org/citations/24318877" target="\_blank">24318877</a>, PubMed:<a href="http://www.uniprot.org/citations/26865365" target="blank">26865365</a>). Maintains protein homeostasis during cellular stress through two opposing mechanisms: protein refolding and degradation. Its acetylation/deacetylation state determines whether it functions in protein refolding or protein degradation by controlling the competitive binding of co-chaperones HOPX and



STUB1. During the early stress response, the acetylated form binds to HOPX which assists in chaperone-mediated protein refolding, thereafter, it is deacetylated and binds to ubiquitin ligase STUB1 that promotes ubiquitin-mediated protein degradation (PubMed:<a href="http://www.uniprot.org/citations/27708256" target="\_blank">27708256</a>). Regulates centrosome integrity during mitosis, and is required for the maintenance of a functional mitotic centrosome that supports the assembly of a bipolar mitotic spindle (PubMed:<a href="http://www.uniprot.org/citations/27137183" target="\_blank">27137183</a>). Enhances STUB1-mediated SMAD3 ubiquitination and degradation and facilitates STUB1-mediated inhibition of TGF-beta signaling (PubMed:<a href="http://www.uniprot.org/citations/24613385" target="\_blank">24613385</a>). Essential for STUB1-mediated ubiquitination and degradation of FOXP3 in regulatory T-cells (Treg) during inflammation (PubMed:<a href="http://www.uniprot.org/citations/23973223" target="\_blank">23973223</a>).

#### **Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs

#### **Tissue Location**


HSPA1B is testis-specific.

### **HSP70 Mouse mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **HSP70 Mouse mAb - Images**




Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP70) Monoclonal Antibody, Unconjugated (AP94160) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.






Sample: Hela Cell (Human) Lysate at 40 ug Primary: Anti-HSP70 (AP94160) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 70 kD




Sample: HepG2 Cell (Human) Lysate at 40 ug Primary: Anti-HSP70 (AP94160) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 70 kD




Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium



citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP70) Monoclonal Antibody, Unconjugated (ascites of AP94160 5G10) at 1:2000 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP70) Monoclonal Antibody, Unconjugated (ascites of AP94160 5G10) at 1:2000 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP70) Monoclonal Antibody, Unconjugated (ascites of AP94160 5G10) at 1:2000 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

## **HSP70 Mouse mAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.