

Galectin 9 Rabbit pAb
Galectin 9 Rabbit pAb
Catalog # AP94163**Specification**

Galectin 9 Rabbit pAb - Product Information

Application	IHC-P
Primary Accession	O08573
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40036

Galectin 9 Rabbit pAb - Additional Information**Gene ID** 16859**Other Names**

Galectin-9, Gal-9, Lgals9

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Galectin 9 Rabbit pAb - Protein Information**Name** Lgals9**Function**

Binds galactosides (By similarity). Has high affinity for the Forssman pentasaccharide (By similarity). Ligand for HAVCR2/TIM3 (By similarity). Binding to HAVCR2 induces T-helper type 1 lymphocyte (Th1) death (By similarity). Also stimulates bactericidal activity in infected macrophages by causing macrophage activation and IL1B secretion which restricts intracellular bacterial growth (PubMed:20937702). Ligand for P4HB; the interaction retains P4HB at the cell surface of Th2 T-helper cells, increasing disulfide reductase activity at the plasma membrane, altering the plasma membrane redox state and enhancing cell migration (PubMed:21670307). Ligand for CD44; the interaction enhances binding of SMAD3 to the FOXP3 promoter, leading to up-regulation of FOXP3 expression and increased induced regulatory T (iTreg) cell stability and suppressive function (PubMed:25065622). Promotes ability of mesenchymal stromal cells to suppress T-cell proliferation (By similarity). Expands regulatory T- cells and induces cytotoxic T-cell apoptosis following virus infection (By similarity). Activates ERK1/2 phosphorylation inducing cytokine (IL-6, IL-8, IL-12) and chemokine (CCL2) production in mast and dendritic cells (By

similarity). Inhibits degranulation and induces apoptosis of mast cells (By similarity). Induces maturation and migration of dendritic cells (By similarity). Inhibits natural killer (NK) cell function (PubMed:23408620). Can transform NK cell phenotype from peripheral to decidual during pregnancy (By similarity). Astrocyte derived galectin-9 enhances microglial TNF production (PubMed:25158758). May play a role in thymocyte-epithelial interactions relevant to the biology of the thymus. May provide the molecular basis for urate flux across cell membranes, allowing urate that is formed during purine metabolism to efflux from cells and serving as an electrogenic transporter that plays an important role in renal and gastrointestinal urate excretion (By similarity). Highly selective to the anion urate (By similarity).

Cellular Location

Cytoplasm. Nucleus. Secreted {ECO:0000250|UniProtKB:O00182} Note=May also be secreted by a non-classical secretory pathway (PubMed:9038233). Secreted by mesenchymal stromal cells upon IFNG stimulation (By similarity). {ECO:0000250|UniProtKB:O00182, ECO:0000269|PubMed:9038233}

Tissue Location

Accentuated expression in liver and thymus of embryo, detected in embryonic heart, brain, lung, liver, and kidney Highly expressed in adult thymus, small intestine, and liver, and to a lesser extent in lung, kidney, spleen, cardiac, and skeletal muscle Barely detectable in brain and reticulocyte. Expressed in placenta, uterus and decidua during pregnancy (PubMed:23242525). Expressed in CD4+ T-cells with higher levels in iTreg cells than other T-cell types and sustained high levels throughout iTreg cell differentiation (at protein level) (PubMed:25065622). Expressed in myeloid cells in lung (PubMed:20937702). Constitutively expressed in microglia (PubMed:25158758). Isoform 1 is expressed exclusively in the small intestine. Isoform 2 expression in decidua increases in pathological pregnancy from gestation day 7.5 to 13.5 and it is higher than in normal pregnancy (PubMed:23242525). Isoform 3 expression in decidua is higher in normal pregnancy than in pathological pregnancy (PubMed:23242525).


Galectin 9 Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Galectin 9 Rabbit pAb - Images







Tissue/cell: rat liver tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Galectin-9 Polyclonal Antibody, Unconjugated(AP94163) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Tissue/cell: mouse kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Galectin-9 Polyclonal Antibody, Unconjugated(AP94163) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (AP94163) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (AP94163) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (AP94163) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human tonsil); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (galectin 9) Polyclonal Antibody, Unconjugated (AP94163) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Galectin 9 Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.