

Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb
Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb
Catalog # AP94264**Specification**

Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Host	Rabbit
Clonality	Recombinant

Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Additional Information**Dilution**

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Protein Information**Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Images



Cell line: HeLa Fixative: 4% Paraformaldehyde Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Mouse IgG Nuclear counter stain: DAPI (Blue) Counter stain: Tubulin (Red) Comment: Color green is the positive signal for AP94264




Tissue: Rat stomach Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Mouse)(sp-0024) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for AP94264




Tissue: Mouse kidney Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Mouse)(sp-0024) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive

signal for AP94264



Tissue: Human liver Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Mouse)(sp-0024) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for AP94264



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 4°C overnight Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate: 1: HeLa, 2: Neuro-2a, 3: BRL Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 15 kDa Observed MW: 15 kDa

Di-Methyl-Histone H3 (Lys27) Recombinant Mouse mAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.