

Butyryl-Histone H2B (Lys20) Rabbit pAb
Butyryl-Histone H2B (Lys20) Rabbit pAb
Catalog # AP94501**Specification**

Butyryl-Histone H2B (Lys20) Rabbit pAb - Product Information

Application	WB, IF, ICC
Host	Rabbit
Clonality	Polyclonal

Butyryl-Histone H2B (Lys20) Rabbit pAb - Additional Information**Dilution**

WB ~ 1:1000
IF ~ 1:50 ~ 200
ICC ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Butyryl-Histone H2B (Lys20) Rabbit pAb - Protein Information**Butyryl-Histone H2B (Lys20) Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Butyryl-Histone H2B (Lys20) Rabbit pAb - Images



Cell line: (A) HeLa, (B) HeLa+SBA (5mM, 24hr) Fixative: 100% Ice-cold methanol
Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Counter stain: Tubulin (Red) Comment: Color green is the positive signal for AP94501



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 14 kDa Observed MW: 14 kDa

Butyryl-Histone H2B (Lys20) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.