

phospho-PPAR alpha (Ser12) Rabbit pAb
phospho-PPAR alpha (Ser12) Rabbit pAb
Catalog # AP94506

Specification

phospho-PPAR alpha (Ser12) Rabbit pAb - Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q6I9S0
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

phospho-PPAR alpha (Ser12) Rabbit pAb - Additional Information

Dilution

WB~1:1000
IHC-P~N/A
IHC-F~N/A
IF~1:50~200
E~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


phospho-PPAR alpha (Ser12) Rabbit pAb - Protein Information


phospho-PPAR alpha (Ser12) Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

phospho-PPAR alpha (Ser12) Rabbit pAb - Images






phosphopeptide non phosphopeptide



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-phospho-PPAR alpha(Ser12) Polyclonal Antibody, Unconjugated(AP94506) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample: Lane 1: Heart (Mouse) Lysate at 40 ug Lane 2: Liver (Mouse) Lysate at 40 ug Lane 3: Kidney (Mouse) Lysate at 40 ug Lane 4: Cerebrum (Mouse) Lysate at 40 ug Lane 5: Cerebrum (Rat) Lysate at 40 ug Lane 6: Jurkat (Human) Cell Lysate at 30 ug Lane 7: A431 (Human) Cell Lysate at 30 ug Primary: Anti-phospho-PPAR alpha (Ser12) (AP94506) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 53 kD Observed band size: 50 kD

phospho-PPAR alpha (Ser12) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.