

**Acetyl-Histone H3 (Lys27) Rabbit pAb**  
**Acetyl-Histone H3 (Lys27) Rabbit pAb**  
**Catalog # AP94605****Specification**

---

**Acetyl-Histone H3 (Lys27) Rabbit pAb - Product Information**

|             |            |
|-------------|------------|
| Application | IHC-P, WB  |
| Host        | Rabbit     |
| Clonality   | Polyclonal |

**Acetyl-Histone H3 (Lys27) Rabbit pAb - Additional Information****Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


**Storage**


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Acetyl-Histone H3 (Lys27) Rabbit pAb - Protein Information****Acetyl-Histone H3 (Lys27) Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Acetyl-Histone H3 (Lys27) Rabbit pAb - Images**



Tissue: Human breast cancer Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:200 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for AP94605



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:400 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: Pharbitis, N2a, Mouse kidney Protein loading quantity: 20 µg Exposure time: 30 s Predicted MW: 17 kDa Observed MW: 17 kDa



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 17 kDa Observed MW: 17 kDa

**Acetyl-Histone H3 (Lys27) Rabbit pAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.