

Phospho-Histone H3 (Thr45) Rabbit pAb
Phospho-Histone H3 (Thr45) Rabbit pAb
Catalog # AP94670**Specification**

Phospho-Histone H3 (Thr45) Rabbit pAb - Product Information

Application	WB
Host	Rabbit
Clonality	Polyclonal

Phospho-Histone H3 (Thr45) Rabbit pAb - Additional Information**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Phospho-Histone H3 (Thr45) Rabbit pAb - Protein Information**Phospho-Histone H3 (Thr45) Rabbit pAb - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Histone H3 (Thr45) Rabbit pAb - Images



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) JK, (+) JK+ TPA (200nM, 30min) + Calyculin A (100nM, 1hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 17 kDa Observed MW: 17 kDa



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+ Nocodazole (100ng/ml, 18hr) + Calyculin A (100nM, 1hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 17 kDa Observed MW: 17 kDa

Phospho-Histone H3 (Thr45) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.