

CYP7A1 Rabbit pAb
CYP7A1 Rabbit pAb
Catalog # AP94705**Specification****CYP7A1 Rabbit pAb - Product Information**

Application	WB
Primary Accession	Q64505
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse CYP7A1
Epitope Specificity	50-150/504
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein. Belongs to the cytochrome P450 family.
SIMILARITY	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Important Note	

Background Descriptions

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first reaction in the cholesterol catabolic pathway in the liver, which converts cholesterol to bile acids. This reaction is the rate limiting step and the major site of regulation of bile acid synthesis, which is the primary mechanism for the removal of cholesterol from the body. Polymorphisms in the promoter of this gene are associated with defects in bile acid synthesis. [provided by RefSeq, Feb 2010].

CYP7A1 Rabbit pAb - Additional Information**Gene ID** 13122**Other Names**

Cytochrome P450 7A1, Cyp7a1 {ECO:0000303|PubMed:14522988, ECO:0000312|MGI:MGI:106091}

Target/Specificity

Detected in liver.

Dilution

WB~~1:1000

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CYP7A1 Rabbit pAb - Protein Information

Name Cyp7a1 {ECO:0000303|PubMed:14522988, ECO:0000312|MGI:MGI:106091}

Function

A cytochrome P450 monooxygenase involved in the metabolism of endogenous cholesterol and its oxygenated derivatives (oxysterols) (By similarity). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (By similarity). Functions as a critical regulatory enzyme of bile acid biosynthesis and cholesterol homeostasis. Catalyzes the hydroxylation of carbon hydrogen bond at 7-alpha position of cholesterol, a rate-limiting step in cholesterol catabolism and bile acid biosynthesis (Probable). 7-alpha hydroxylates several oxysterols, including 4beta-hydroxycholesterol and 24-hydroxycholesterol. Catalyzes the oxidation of the 7,8 double bond of 7-dehydrocholesterol and lathosterol with direct and predominant formation of the 7-keto derivatives (By similarity).


Cellular Location


Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P22680}; Single-pass membrane protein {ECO:0000250|UniProtKB:P22680}. Microsome membrane {ECO:0000250|UniProtKB:P22680}; Single-pass membrane protein {ECO:0000250|UniProtKB:P22680}

CYP7A1 Rabbit pAb - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CYP7A1 Rabbit pAb - Images



Sample: HepG2(Human) Cell Lysate at 30 ug Primary: Anti- CYP7A1 (AP94705) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kD
Observed band size: 55 kD



Sample: Liver (Mouse) Lysate at 40 ug Primary: Anti-CYP7A1 (AP94705) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kD
Observed band size: 55 kD

CYP7A1 Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.