

Glucose 6 phosphate isomerase Recombinant Rabbit mAb
Glucose 6 phosphate isomerase Recombinant Rabbit mAb
Catalog # AP94817

Specification

Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Product Information

Application	WB, IHC-P
Host	Rabbit
Clonality	Recombinant

Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Additional Information

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Protein Information


Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Images







25 ug total protein per lane of various lysates (see on figure) probed with Glucose 6 phosphate isomerase monoclonal antibody, unconjugated (AP94817) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Colon ; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Glucose 6 phosphate isomerase Monoclonal Antibody, Unconjugated(AP94817) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.

Glucose 6 phosphate isomerase Recombinant Rabbit mAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.