

HBsAg Goat pAb
HBsAg Goat pAb
Catalog # AP94833**Specification**

HBsAg Goat pAb - Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q89714
Reactivity	Human
Host	Goat
Clonality	Polyclonal

HBsAg Goat pAb - Additional Information**Dilution**

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HBsAg Goat pAb - Protein Information**HBsAg Goat pAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HBsAg Goat pAb - Images



Paraformaldehyde-fixed, paraffin embedded (Hepatitis B virus infected human hepatocellular carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HBsAg) Polyclonal Antibody, Unconjugated (bs-1557G) at 1:200 overnight at 4°C, followed by A594 conjugated Rabbit Anti-Goat IgG H&L secondary antibody (AP94833) incubation and staining.

HBsAg Goat pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.