

PCYT1A Rabbit pAb
PCYT1A Rabbit pAb
Catalog # AP94840

Specification

PCYT1A Rabbit pAb - Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P49585
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PCYT1A
Epitope Specificity	201-300/367
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm; cytosol. Membrane. It can interconvert between an inactive cytosolic form and an active membrane-bound form. Belongs to the cytidylyltransferase family. Homodimer.
SIMILARITY	The serine residues of the C-terminus are phosphorylated. The inactive soluble form is stabilized by phosphorylation, the active membrane bound form is promoted by anionic lipids or diacylglycerol, and is stabilized by dephosphorylation.
SUBUNIT	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Post-translational modifications	
Important Note	

PCYT1A Rabbit pAb - Additional Information

Gene ID 5130

Other Names

Choline-phosphate cytidylyltransferase A, 2.7.7.15, CCT-alpha, CTP:phosphocholine cytidylyltransferase A, CCT A, CT A, Phosphorylcholine transferase A, PCYT1A, CTPCT, PCYT1

Target/Specificity

Cancer, Neurotrophins, Lipid metabolism

Dilution

IHC-P~~N/A<br \><span class

=IHC-F~~N/A<br \>IF~~1:50~200

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PCYT1A Rabbit pAb - Protein Information

Name PCYT1A

Synonyms CTPCT, PCYT1

Function

Catalyzes the key rate-limiting step in the CDP-choline pathway for phosphatidylcholine biosynthesis.

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:P19836}. Membrane {ECO:0000250|UniProtKB:P19836}; Peripheral membrane protein {ECO:0000250|UniProtKB:P19836}. Endoplasmic reticulum membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P19836}. Nucleus Note=It can interconvert between an inactive cytosolic form and an active membrane-bound form. {ECO:0000250|UniProtKB:P19836}

Tissue Location


Brain, placenta, liver, fetal and adult lung.

PCYT1A Rabbit pAb - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PCYT1A Rabbit pAb - Images




Paraformaldehyde-fixed, paraffin embedded (rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCYT1A) Polyclonal Antibody, Unconjugated (AP94840) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (human pancreatic cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCYT1A) Polyclonal Antibody, Unconjugated (AP94840) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCYT1A) Polyclonal Antibody, Unconjugated (AP94840) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human stomach carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCYT1A) Polyclonal Antibody, Unconjugated (AP94840) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCYT1A) Polyclonal Antibody, Unconjugated (AP94840) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

PCYT1A Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.