

RBP2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9557b

Specification

RBP2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P50120
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15707
Antigen Region	56-84

RBP2 Antibody (Center) - Additional Information

Gene ID 5948

Other Names

Retinol-binding protein 2, Cellular retinol-binding protein II, CRBP-II, RBP2, CRBP2

Target/Specificity

This RBP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 56-84 amino acids from the Central region of human RBP2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RBP2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

RBP2 Antibody (Center) - Protein Information

Name RBP2

Synonyms CRBP2

Function Intracellular transport of retinol.

Cellular Location

Cytoplasm.

Tissue Location


Higher expression in adult small intestine and to a much lesser extent in fetal kidney.

RBP2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBP2 Antibody (Center) - Images



Western blot analysis of RBP2 Antibody (Center) (Cat. #AP9557b) in 293 cell line lysates (35ug/lane). RBP2 (arrow) was detected using the purified Pab.

RBP2 Antibody (Center) - Background

RBP2 is an abundant protein present in the small intestinal epithelium. It is thought to participate in the uptake and/or intracellular metabolism of vitamin A. Vitamin A is a fat-soluble vitamin necessary for growth, reproduction, differentiation of epithelial tissues, and vision. RBP2 may also modulate the supply of retinoic acid to the nuclei of endometrial cells during the menstrual cycle.

RBP2 Antibody (Center) - References

Zeng, J., et al. Gastroenterology 138(3):981-992(2010)

Yamaguchi, N., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 296 (3), G524-G533 (2009) :

?Tarter, M., et al. Proteins 70(4):1626-1630(2008)
?Zhang, L., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 282 (6), G1079-G1087 (2002) :
?Folli, C., et al. Proc. Natl. Acad. Sci. U.S.A. 98(7):3710-3715(2001)
?Herr, F.M., et al. Biochemistry 31(29):6748-6755(1992)
?Astrom, A., et al. J. Biol. Chem. 266(26):17662-17666(1991)