

CBA1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9682a**Specification**

CBA1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9BPX6
Other Accession	Q4R518
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	54351
Antigen Region	52-81

CBA1 Antibody (N-term) - Additional Information**Gene ID** 10367**Other Names**

Calcium uptake protein 1, mitochondrial, Atopy-related autoantigen CALC, ara CALC,
Calcium-binding atopy-related autoantigen 1, Hom s 4, MICU1, CALC, CBARA1

Target/Specificity

This CBA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 52-81 amino acids from the N-terminal region of human CBA1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CBA1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CBA1 Antibody (N-term) - Protein Information**Name** MICU1 {ECO:0000303|PubMed:20693986, ECO:0000312|HGNC:HGNC:1530}

Function Calcium sensor of the mitochondrial calcium uniporter (MCU) channel, which senses calcium level via its EF-hand domains (PubMed:[20693986](#), PubMed:[23101630](#), PubMed:[23747253](#), PubMed:[24313810](#), PubMed:[24332854](#), PubMed:[24503055](#), PubMed:[24560927](#), PubMed:[26341627](#), PubMed:[26903221](#), PubMed:[27099988](#), PubMed:[28615291](#), PubMed:[30454562](#), PubMed:[30638448](#), PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#), PubMed:[32790952](#), PubMed:[34463251](#), PubMed:[36206740](#), PubMed:[37036971](#), PubMed:[37126688](#)). MICU1 and MICU2 (or MICU3) form a disulfide-linked heterodimer that stimulates and inhibits MCU activity, depending on the concentration of calcium (PubMed:[24560927](#), PubMed:[26903221](#), PubMed:[28615291](#), PubMed:[32148862](#), PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#), PubMed:[32790952](#), PubMed:[36206740](#), PubMed:[37036971](#), PubMed:[37126688](#)). At low calcium levels, MICU1 occludes the pore of the MCU channel, preventing mitochondrial calcium uptake (PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#), PubMed:[37036971](#), PubMed:[37126688](#)). At higher calcium levels, calcium-binding to MICU1 and MICU2 (or MICU3) induces a conformational change that weakens MCU-MICU1 interactions and moves the MICU1-MICU2 heterodimer away from the pore, allowing calcium permeation through the MCU channel (PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#)). Also required to protect against manganese toxicity by preventing manganese uptake by MCU: mechanistically, manganese-binding to its EF-hand domains does not induce any conformational change, maintaining MCU pore occlusion (PubMed:[30082385](#), PubMed:[30403999](#)). Also acts as a barrier for inhibitors of the MCU channel, such as ruthenium red or its derivative Ru360 (PubMed:[37244260](#)). Acts as a regulator of mitochondrial cristae structure independently of its ability to regulate the mitochondrial calcium uniporter channel (PubMed:[31427612](#), PubMed:[37098122](#)). Regulates glucose-dependent insulin secretion in pancreatic beta-cells by regulating mitochondrial calcium uptake (PubMed:[22904319](#)). Induces T-helper 1-mediated autoreactivity, which is accompanied by the release of IFNG (PubMed:[16002733](#)).

Cellular Location

Mitochondrion intermembrane space. Mitochondrion inner membrane. Note=Recruited to the mitochondrial inner membrane by EMRE/SMDT1 (PubMed:[30454562](#)). Also localizes to mitochondrial cristae junctions (PubMed:[31427612](#))

Tissue Location


Expressed in epithelial cell lines. Strongly expressed in epidermal keratinocytes and dermal endothelial cells


CBAA1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CBAA1 Antibody (N-term) - Images





Western blot analysis of CBA1 Antibody (N-term) (Cat. #AP9682a) in K562 cell line lysates (35ug/lane). CBA1 (arrow) was detected using the purified Pab.

CBA1 Antibody (N-term) - References

Bordicchia, M., et al. Metab. Clin. Exp. (2009) In press :
Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)
Natter, S., et al. FASEB J. 12(14):1559-1569(1998)