

DGAT2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9686a

Specification

DGAT2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q96PD7
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

DGAT2 Antibody (N-term) - Additional Information

Gene ID 84649

Other Names

Diacylglycerol O-acyltransferase 2, Acyl-CoA retinol O-fatty-acyltransferase, ARAT, Retinol O-fatty-acyltransferase, Diglyceride acyltransferase 2, DGAT2

Target/Specificity

This DGAT2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human DGAT2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DGAT2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DGAT2 Antibody (N-term) - Protein Information

Name DGAT2 ([HGNC:16940](#))

Function Essential acyltransferase that catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates. Required for

synthesis and storage of intracellular triglycerides (PubMed:[27184406](#)). Probably plays a central role in cytosolic lipid accumulation. In liver, is primarily responsible for incorporating endogenously synthesized fatty acids into triglycerides (By similarity). Also functions as an acyl-CoA retinol acyltransferase (ARAT) (By similarity). Also able to use 1- monoalkylglycerol (1-MAKG) as an acyl acceptor for the synthesis of monoalkyl-monoacylglycerol (MAMAG) (PubMed:[28420705](#)).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Lipid droplet. Cytoplasm, perinuclear region

Tissue Location


Predominantly expressed in liver and white adipose tissue. Expressed at lower level in mammary gland, testis and peripheral blood leukocytes. Expressed in sebaceous glands of normal skin but decreased psoriatic skin.

DGAT2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DGAT2 Antibody (N-term) - Images



Western blot analysis of DGAT2 Antibody (N-term) (Cat. #AP9686a) in NCI-H460 cell line lysates (35ug/lane). DGAT2 (arrow) was detected using the purified Pab.

DGAT2 Antibody (N-term) - Background

Acyl-CoA:diacylglycerol acyltransferase, or DGAT (EC 2.3.1.20), is responsible for the synthesis of triglycerides. It catalyzes a reaction in which diacylglycerol is covalently joined to long chain fatty

acyl-CoAs.

DGAT2 Antibody (N-term) - References

- # Kantartzis, K., et al. Clin. Sci. 116(6):531-537(2009)
- # Stone, S.J., et al. J. Biol. Chem. 284(8):5352-5361(2009)
- # Yen, C.L., et al. J. Lipid Res. 49(11):2283-2301(2008)
- # Levin, M.C., et al. Am. J. Physiol. Endocrinol. Metab. 293 (6), E1772-E1781 (2007)
- # Payne, V.A., et al. J. Biol. Chem. 282(29):21005-21014(2007)
- # Yamada, S., et al. Oncogene 23(35):5901-5911(2004)
- # Wakimoto, K., et al. Biochem. Biophys. Res. Commun. 310(2):296-302(2003)