

KRT1 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9695c**Specification**

KRT1 Antibody (Center) - Product Information

Application	WB, FC, IHC-P,E
Primary Accession	P04264
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	415-443

KRT1 Antibody (Center) - Additional Information**Gene ID** 3848**Other Names**

Keratin, type II cytoskeletal 1, 67 kDa cytokeratin, Cytokeratin-1, CK-1, Hair alpha protein, Keratin-1, K1, Type-II keratin Kb1, KRT1, KRTA

Target/Specificity

This KRT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 415-443 amino acids from the Central region of human KRT1.

Dilution

WB~~1:1000
FC~~1:10~50
IHC-P~~1:50~100
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KRT1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KRT1 Antibody (Center) - Protein Information**Name** KRT1

Synonyms KRTA

Function May regulate the activity of kinases such as PKC and SRC via binding to integrin beta-1 (ITB1) and the receptor of activated protein C kinase 1 (RACK1). In complex with C1QBP is a high affinity receptor for kininogen-1/HMWK.

Cellular Location

Cell membrane. Cytoplasm

Tissue Location


The source of this protein is neonatal foreskin. The 67-kDa type II keratins are expressed in terminally differentiating epidermis

KRT1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KRT1 Antibody (Center) - Images




All lanes: Anti-KRT1 Antibody (Center) at 1:1000 dilution + 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 66KDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of KRT1 Antibody (Center) (Cat. #AP9695c) in MCF-7 cell line lysates (35ug/lane). KRT1 (arrow) was detected using the purified Pab.



KRT1 Antibody (Center) (Cat. #AP9695c) IHC analysis in formalin fixed and paraffin embedded skin tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the KRT1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



KRT1 Antibody (Center) (Cat. #AP9695c) flow cytometric analysis of MCF-7 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

KRT1 Antibody (Center) - Background

KRT1 is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically

expressed in the spinous and granular layers of the epidermis with family member KRT10 and mutations in these genes have been associated with bullous congenital ichthyosiform erythroderma.

KRT1 Antibody (Center) - References

Labudova, M., et al. J. Virol. 83(16):7842-7849(2009)
Barcelos, A.C., et al. J. Cutan. Pathol. 36(6):647-654(2009)
Grimberg, G., et al. Br. J. Dermatol. 160(2):446-449(2009)

KRT1 Antibody (Center) - Citations

- [Differential miRNA expression profiles in human keratinocytes in response to protein kinase C inhibitor.](#)