

ADPRHL2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9723a

Specification

ADPRHL2 Antibody (N-term) - Product Information

Application	FC, WB,E
Primary Accession	Q9NX46
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	87-114

ADPRHL2 Antibody (N-term) - Additional Information

Gene ID 54936

Other Names

Poly(ADP-ribose) glycohydrolase ARH3, ADP-ribosylhydrolase 3, [Protein ADP-ribosylarginine] hydrolase-like protein 2, ADPRHL2, ARH3

Target/Specificity

This ADPRHL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 87-114 amino acids from the N-terminal region of human ADPRHL2.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ADPRHL2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ADPRHL2 Antibody (N-term) - Protein Information

Name ADPRS ([HGNC:21304](#))

Function ADP-ribosylhydrolase that preferentially hydrolyzes the scissile alpha-O-linkage attached

to the anomeric C1'' position of ADP- ribose and acts on different substrates, such as proteins ADP-ribosylated on serine and threonine, free poly(ADP-ribose) and O- acetyl-ADP-D-ribose (PubMed:[21498885](#), PubMed:[29907568](#), PubMed:[30045870](#), PubMed:[30401461](#), PubMed:[30830864](#), PubMed:[33186521](#), PubMed:[33769608](#), PubMed:[33894202](#), PubMed:[34019811](#), PubMed:[34321462](#), PubMed:[34479984](#), PubMed:[34625544](#)). Specifically acts as a serine mono-ADP- ribosylhydrolase by mediating the removal of mono-ADP-ribose attached to serine residues on proteins, thereby playing a key role in DNA damage response (PubMed:[28650317](#), PubMed:[29234005](#), PubMed:[30045870](#), PubMed:[33186521](#), PubMed:[34019811](#), PubMed:[34625544](#)). Serine ADP- ribosylation of proteins constitutes the primary form of ADP- ribosylation of proteins in response to DNA damage (PubMed:[29480802](#), PubMed:[33186521](#), PubMed:[34625544](#)). Does not hydrolyze ADP-ribosyl- arginine, -cysteine, -diphthamide, or -asparagine bonds (PubMed:[16278211](#), PubMed:[33769608](#)). Also able to degrade protein free poly(ADP-ribose), which is synthesized in response to DNA damage: free poly(ADP-ribose) acts as a potent cell death signal and its degradation by ADPRHL2 protects cells from poly(ADP-ribose)-dependent cell death, a process named parthanatos (PubMed:[16278211](#)). Also hydrolyzes free poly(ADP-ribose) in mitochondria (PubMed:[22433848](#)). Specifically digests O-acetyl-ADP-D-ribose, a product of deacetylation reactions catalyzed by sirtuins (PubMed:[17075046](#), PubMed:[21498885](#)). Specifically degrades 1''-O-acetyl-ADP-D-ribose isomer, rather than 2''-O-acetyl- ADP-D-ribose or 3''-O-acetyl-ADP-D-ribose isomers (PubMed:[21498885](#)).

Cellular Location

Nucleus. Cytoplasm. Chromosome Mitochondrion matrix Note=Recruited to DNA lesion regions following DNA damage; ADP-D- ribose-recognition is required for recruitment to DNA damage sites

Tissue Location

Ubiquitous (PubMed:[16278211](#)). Expressed in skin fibroblasts (PubMed:[30830864](#)).


ADPRHL2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADPRHL2 Antibody (N-term) - Images





Western blot analysis of ADPRHL2 Antibody (N-term) (Cat. #AP9723a) in HeLa cell line lysates (35ug/lane). ADPRHL2 (arrow) was detected using the purified Pab.



ADPRHL2 Antibody (N-term) (Cat. #AP9723a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADPRHL2 Antibody (N-term) - Background

ADPRHL2 is a member of the ADP-ribosylglycohydrolase family. The enzyme catalyzes the removal of ADP-ribose from ADP-ribosylated proteins. This enzyme localizes to the mitochondria, in addition to the nucleus and cytoplasm.

ADPRHL2 Antibody (N-term) - References

- Niere, M., et al. Mol. Cell. Biol. 28(2):814-824(2008)
- Ono, T., et al. Proc. Natl. Acad. Sci. U.S.A. 103(45):16687-16691(2006)
- Mueller-Dieckmann, C., et al. Proc. Natl. Acad. Sci. U.S.A. 103(41):15026-15031(2006)