

### **RPS24 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9726C

### Specification

# **RPS24 Antibody (Center) - Product Information**

Application Primary Accession Other Accession

Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region FC, WB,E <u>P62847</u> <u>P62850, P62849, Q4R5H5, Q56JU9,</u> <u>A0A5F9D2E6</u> Human Bovine, Monkey, Mouse, Rabbit, Rat Rabbit Polyclonal Rabbit IgG 15423 65-93

## **RPS24** Antibody (Center) - Additional Information

Gene ID 6229

Other Names 40S ribosomal protein S24, RPS24

#### Target/Specificity

This RPS24 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 65-93 amino acids from the Central region of human RPS24.

**Dilution** FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

RPS24 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## **RPS24 Antibody (Center) - Protein Information**



Name RPS24 (HGNC:10411)

**Function** Component of the small ribosomal subunit (PubMed:<u>23636399</u>). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:<u>23636399</u>). Required for processing of pre-rRNA and maturation of 40S ribosomal subunits (PubMed:<u>18230666</u>). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:<u>34516797</u>).

Cellular Location Cytoplasm. Nucleus, nucleolus

#### **Tissue Location**


Mature tissues, such as adult brain, skeletal muscle, heart, and kidney, express low levels, whereas tissues and organs with significant populations of proliferating cells, such as fetal brain, placenta, bone marrow, and various glandular organs, contain significantly higher levels.

## **RPS24 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


### **RPS24 Antibody (Center) - Images**



Anti-RPS24 Antibody (Center) at 1:1000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.



Predicted band size : 15 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



RPS24 Antibody (Center) (Cat. #AP9726c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# **RPS24 Antibody (Center) - Background**

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. RPS24 is a ribosomal protein that is a component of the 40S subunit. The protein belongs to the S24E family of ribosomal proteins. It is located in the cytoplasm. Multiple transcript variants encoding different isoforms have been found for this gene. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

## **RPS24 Antibody (Center) - References**

Quarello, P., et al. Haematologica 95(2):206-213(2010) Badhai, J., et al. Biochim. Biophys. Acta 1792(10):1036-1042(2009) Robledo, S., et al. RNA 14(9):1918-1929(2008)