10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MAPK1IP1L Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9730c

Specification

MAPK1IP1L Antibody (Center) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	Q8NDC0
Other Accession	Q4R837, Q5E9L3
Reactivity	**Human**
Predicted	**Bovine, Monkey**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**24269**
Antigen Region	**108-137**

MAPK1IP1L Antibody (Center) - Additional Information

Gene ID 93487

Other Names
MAPK-interacting and spindle-stabilizing protein-like, Mitogen-activated protein kinase 1-interacting protein 1-like, MAPK1IP1L, C14orf32

Target/Specificity
This MAPK1IP1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 108-137 amino acids from the Central region of human MAPK1IP1L.

Dilution
WB~~1:1000
IHC-P~~1:50~100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MAPK1IP1L Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MAPK1IP1L Antibody (Center) - Protein Information

Name MAPK1IP1L

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Synonyms C14orf32

MAPK1IP1L Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MAPK1IP1L Antibody (Center) - Images



All lanes : Anti-MAPK1IP1L Antibody (Center) at 1:1000 dilution Lane 1: U-251 MG whole cell lysate Lane 2: Y79 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



MAPK1IP1L Antibody (Center) (Cat. #AP9730c) IHC analysis in formalin fixed and paraffin embedded kidney carcinoma followed by peroxidase conjugation of the secondary antibody and

DAB staining. This data demonstrates the use of the MAPK1IP1L Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

MAPK1IP1L Antibody (Center) - References

?Olsen, J.V., et al. Cell 127(3):635-648(2006)
?Heilig, R., et al. Nature 421(6923):601-607(2003)