

KLC2 Antibody (C-Term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9798b**Specification**

KLC2 Antibody (C-Term) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q9H0B6
Other Accession	NP_073733
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	403-432

KLC2 Antibody (C-Term) - Additional Information**Gene ID** 64837**Other Names**

Kinesin light chain 2, KLC 2, KLC2

Target/Specificity

This KLC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 403-432 amino acids from the C-terminal region of human KLC2.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KLC2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

KLC2 Antibody (C-Term) - Protein Information**Name** KLC2 ([HGNC:20716](#))

Function Kinesin is a microtubule-associated force-producing protein that plays a role in organelle transport. The light chain functions in coupling of cargo to the heavy chain or in the modulation of its ATPase activity (Probable). Through binding with PLEKHM2 and ARL8B, recruits kinesin-1 to lysosomes and hence direct lysosomes movement toward microtubule plus ends (PubMed:[22172677](#)).

Cellular Location


Cytoplasm, cytoskeleton. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side

KLC2 Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KLC2 Antibody (C-Term) - Images



Western blot analysis of KLC2 Antibody (C-Term) (Cat. #AP9798b) in mouse NIH-3T3 cell line lysates (35ug/lane). KLC2 (arrow) was detected using the purified Pab.



KLC2 Antibody (C-Term) (Cat. #AP9798b) IHC analysis in formalin fixed and paraffin embedded brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the KLC2 Antibody (C-Term) for immunohistochemistry. Clinical relevance has not been evaluated.



KLC2 Antibody (C-Term) (Cat. #AP9798b) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

KLC2 Antibody (C-Term) - Background

Kinesin is a molecular motor that generates ATP-dependent movement of vesicles and organelles along microtubules. Kinesin consists of 2 light chains, such as KLC2, and 2 heavy chains (see KIF5B; MIM 602809) in a 1:1 stoichiometric ratio.

KLC2 Antibody (C-Term) - References

- Trejo, H.E., et al. FASEB J. 24(2):374-382(2010)
- Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
- Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :
- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)
- Lim, J., et al. Cell 125(4):801-814(2006)