

**DNAJC2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP9885c****Specification**

---

**DNAJC2 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q99543</a>
Other Accession	<a href="#">Q4R8H2</a> , <a href="#">Q1RMH9</a>
Reactivity	Human, Mouse
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	71996
Antigen Region	226-255

**DNAJC2 Antibody (Center) - Additional Information****Gene ID** 27000**Other Names**

Dnaj homolog subfamily C member 2, M-phase phosphoprotein 11, Zuotin-related factor 1, Dnaj homolog subfamily C member 2, N-terminally processed, DNAJC2, MPHOSPH11, MPP11, ZRF1

**Target/Specificity**

This DNAJC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 226-255 amino acids from the Central region of human DNAJC2.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DNAJC2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**DNAJC2 Antibody (Center) - Protein Information****Name** DNAJC2

**Synonyms** MPHOSPH11, MPP11, ZRF1

**Function** Acts both as a chaperone in the cytosol and as a chromatin regulator in the nucleus. When cytosolic, acts as a molecular chaperone: component of the ribosome-associated complex (RAC), a complex involved in folding or maintaining nascent polypeptides in a folding-competent state. In the RAC complex, stimulates the ATPase activity of the ribosome-associated pool of Hsp70-type chaperones HSPA14 that bind to the nascent polypeptide chain. When nuclear, mediates the switching from polycomb-repressed genes to an active state: specifically recruited at histone H2A ubiquitinated at 'Lys-119' (H2AK119ub), and promotes the displacement of the polycomb PRC1 complex from chromatin, thereby facilitating transcription activation.

**Cellular Location**

Nucleus. Cytoplasm, cytosol

**Tissue Location**


Widely expressed..

**DNAJC2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DNAJC2 Antibody (Center) - Images**



Western blot analysis of DNAJC2 Antibody (Center) (Cat. #AP9885c) in mouse Neuro-2a cell line lysates (35ug/lane). DNAJC2 (arrow) was detected using the purified Pab.

**DNAJC2 Antibody (Center) - Background**

This gene is a member of the M-phase phosphoprotein (MPP) family. The gene encodes a phosphoprotein with a J domain and a Myb DNA-binding domain which localizes to both the nucleus and the cytosol. The protein is capable of forming a heterodimeric complex that associates with ribosomes, acting as a molecular chaperone for nascent polypeptide chains as they exit the ribosome. This protein was identified as a leukemia-associated antigen and expression of the gene is upregulated in leukemic blasts. Also, chromosomal aberrations involving this gene are associated with primary head and neck squamous cell tumors. This gene has a pseudogene on chromosome 6.

#### **DNAJC2 Antibody (Center) - References**

Hatzold, J., et al. PLoS Biol. 6 (4), E84 (2008) :  
Olsen, J.V., et al. Cell 127(3):635-648(2006)  
Otto, H., et al. Proc. Natl. Acad. Sci. U.S.A. 102(29):10064-10069(2005)