

**Ki-67**  
**Rabbit Monoclonal Antibody (Mab)**  
**Catalog # APA028**

## Specification

---

### Ki-67 - Product Information

Application	IHC
Primary Accession	<a href="#">P46013</a>
Host	Rabbit
Clonality	Monoclonal
Calculated MW	358694 Da

### Ki-67 - Additional Information

Gene ID	4288
Gene Name	MKI67 ( <a href="#">HGNC:7107</a> )

#### Other Names

Proliferation marker protein Ki-67, Antigen identified by monoclonal antibody Ki-67, Antigen KI-67, Antigen Ki67, MKI67 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=7107](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=7107))

#### Dilution

IHC~~1:100~500

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Ki-67 is for research use only and not for use in diagnostic or therapeutic procedures.

### Ki-67 - Protein Information

**Name** MKI67 ([HGNC:7107](#))

#### Function

Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly (PubMed:[27362226](#)). Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the chromosome surface (PubMed:[27362226](#)). Prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as

**Cellular Location**

a surfactant, dispersing chromosomes and enabling independent chromosome motility (PubMed:[27362226](#)). Binds DNA, with a preference for supercoiled DNA and AT-rich DNA (PubMed:[10878551](#)). Does not contribute to the internal structure of mitotic chromosomes (By similarity). May play a role in chromatin organization (PubMed:[24867636](#)). It is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in maintaining mitotic chromosomes dispersed (Probable).

**Chromosome. Nucleus. Nucleus, nucleolus**  
**Note=**Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the mitotic chromosome surface (PubMed:[27362226](#)). Associates with satellite DNA in G1 phase (PubMed:[9510506](#)). Binds tightly to chromatin in interphase, chromatin-binding decreases in mitosis when it associates with the surface of the condensed chromosomes (PubMed:[15896774](#), PubMed:[22002106](#)). Predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix (PubMed:[22002106](#)).

**Ki-67 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Ki-67 - Images**