

CFTR

Mouse Monoclonal Antibody (Mab)
Catalog # APA156

Specification

CFTR - Product Information

Application IHC
Primary Accession P13569
Host Mouse
Clonality Monoclonal
Calculated MW 168142 Da

CFTR - Additional Information

Gene ID 1080
Gene Name CFTR

Other Names

Cystic fibrosis transmembrane conductance regulator, CFTR, ATP-binding cassette sub-family C member 7, Channel conductance-controlling ATPase, 5.6.1.6, cAMP-dependent chloride channel, CFTR, ABCC7

Storage Maintain refrigerated at 2-8°C for up to 2

weeks. For long term storage store at -20°C in small aliquots to prevent

freeze-thaw cycles.

Precautions CFTR is for research use only and not for

use in diagnostic or therapeutic

procedures.

CFTR - Protein Information

Name CFTR

Synonyms ABCC7

Function Epithelial ion channel that plays an


important role in the regulation of epithelial ion and water transport and fluid

homeostasis (PubMed: 26823428). Mediates the transport of chloride ions across the

cell membrane (PubMed:<u>10792060</u>, PubMed:<u>11524016</u>, PubMed:<u>11707463</u>, PubMed:<u>12519745</u>, PubMed:<u>15010471</u>,

PubMed: 12588899, PubMed: 17036051, PubMed: 19398555, PubMed: 19621064, PubMed: 22178883, PubMed: 25330774,


PubMed: 1712898, PubMed: 8910473, PubMed: 9804160, PubMed: 12529365, PubMed: 17182731, PubMed: 26846474, PubMed: 28087700). Channel activity is





Cellular Location

coupled to ATP hydrolysis (PubMed:8910473). The ion channel is also permeable to HCO(3-); selectivity depends on the extracellular chloride concentration (PubMed: 15010471, PubMed: 19019741). Exerts its function also by modulating the activity of other ion channels and transporters (PubMed: 12403779, PubMed: 22178883, PubMed: 22121115, PubMed: 27941075). Plays an important role in airway fluid homeostasis (PubMed: 16645176, PubMed: 19621064, PubMed: 26823428). Contributes to the regulation of the pH and the ion content of the airway surface fluid layer and thereby plays an important role in defense against pathogens (PubMed: 14668433, PubMed: 16645176, PubMed: 26823428). Modulates the activity of the epithelial sodium channel (ENaC) complex, in part by regulating the cell surface expression of the ENaC complex (PubMed: 17434346, PubMed: 27941075, PubMed: 17182731). Inhibits the activity of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed: 17182731). Inhibits the activity of the ENaC channel containing subunits SCNN1D, SCNN1B and SCNN1G, but not of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed: 27941075). May regulate bicarbonate secretion and salvage in epithelial cells by regulating the transporter SLC4A7 (PubMed: 12403779). Can inhibit the chloride channel activity of ANO1 (PubMed: 22178883). Plays a role in the chloride and bicarbonate homeostasis during sperm epididymal maturation and capacitation (PubMed: 19923167, PubMed:27714810). Apical cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Early endosome membrane: Multi-pass membrane protein {ECO:0000269|Ref.54}. Cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Recycling endosome membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. **Endoplasmic reticulum membrane**; Multi-pass membrane protein {ECO:0000269|Ref.54}. Nucleus {ECO:0000250|UniProtKB:P34158}. Note=The channel is internalized from the cell surface into an endosomal recycling compartment, from where it is recycled to the cell membrane (PubMed:17462998, PubMed:19398555, PubMed:20008117). In





Tissue Location

the oviduct and bronchus, detected on the apical side of epithelial cells, but not associated with cilia (PubMed:22207244). In Sertoli cells, a processed product is detected in the nucleus (By similarity). ER stress induces GORASP2-mediated unconventional (ER/Golgi-independent) trafficking of core-glycosylated CFTR to cell membrane (PubMed:21884936). {ECO:0000250|UniProtKB:P34158, ECO:0000269|PubMed:19398555, ECO:0000269|PubMed:20008117, ECO:0000269|PubMed:21884936, ECO:0000269|PubMed:22207244. ECO:0000305|PubMed:17462998} Expressed in the respiratory airway, including bronchial epithelium, and in the female reproductive tract, including oviduct (at protein level) (PubMed:22207244, PubMed:15716351). **Detected in pancreatic intercalated ducts** in the exocrine tissue, on epithelial cells in intralobular striated ducts in sublingual salivary glands, on apical membranes of crypt cells throughout the small and large intestine, and on the reabsorptive duct in eccrine sweat glands (PubMed:1284548, PubMed:28130590). Detected on the equatorial segment of the sperm head (at protein level) (PubMed:19923167). Detected in nasal and bronchial superficial epithelium (PubMed:15716351). Expressed by the central cells on the sebaceous glands, dermal adipocytes and, at lower levels, by epithelial cells (PubMed:28130590)

CFTR - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CFTR - Images