

BG8,Lewisy
Mouse Monoclonal Antibody (Mab)
Catalog # APA360**Specification****BG8,Lewisy - Product Information**

Application	IHC
Primary Accession	P21217
Host	Mouse
Clonality	Monoclonal
Calculated MW	42117 Da

BG8,Lewisy - Additional Information

Gene ID	2525
Gene Name	FUT3 (HGNC:4014)

Other Names

3-galactosyl-N-acetylglucosaminide 4-alpha-L-fucosyltransferase FUT3, 2.4.1.65,
Alpha-3-fucosyltransferase FUT3, 2.4.1.-, Blood group Lewis alpha-4-fucosyltransferase, Lewis FT,
Fucosyltransferase 3, Fucosyltransferase III, FucT-III, FUT3 (HGNC:4014), FT3B, LE

Dilution

IHC~~1:100~500

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BG8,Lewisy is for research use only and not for use in diagnostic or therapeutic procedures.

BG8,Lewisy - Protein Information**Name** [FUT3 \(HGNC:4014\)](#)**Synonyms****Function****FT3B, LE**

Catalyzes the transfer of L-fucose, from a guanosine diphosphate-beta-L-fucose, to both the subterminal N-acetyl glucosamine (GlcNAc) of type 1 chain (beta-D-Gal-(1->3)-beta-D-GlcNAc) glycolipids and oligosaccharides via an alpha(1,4) linkage, and the subterminal glucose (Glc) or GlcNAc of type 2 chain (beta-D-Gal-(1->4)-beta-D- GlcNAc) oligosaccharides via an alpha(1,3) linkage,

independently of the presence of terminal alpha-L-fucosyl-(1,2) moieties on the terminal galactose of these acceptors and participates in the blood groups Lewis determination and expression of Lewis a (Le(a)), lewis b (Le(b)), Lewis x/SSEA-1 (Le(x)) and lewis y (Le(y)) antigens (PubMed:[12668675](#), PubMed:[1977660](#), PubMed:[11058871](#)). Also catalyzes the transfer of L- fucose to subterminal GlcNAc of sialyl- and disialyl-lactotetraosylceramide to produce sialyl Lewis a (sLe(a)) and disialyl Lewis a via an alpha(1,4) linkage and therefore may regulate cell surface sialyl Lewis a expression and consequently regulates adhesive properties to E-selectin, cell proliferation and migration (PubMed:[12668675](#), PubMed:[11058871](#), PubMed:[27453266](#)). Catalyzes the transfer of an L-fucose to 3'-sialyl-N-acetyllactosamine by an alpha(1,3) linkage, which allows the formation of sialyl-Lewis x structure and therefore may regulate the sialyl-Lewis x surface antigen expression and consequently adhesive properties to E-selectin (PubMed:[11058871](#)). Prefers type 1 chain over type 2 acceptors (PubMed:[7721776](#)). Type 1 tetrasaccharide is a better acceptor than type 1 disaccharide suggesting that a beta anomeric configuration of GlcNAc in the substrate is preferred (PubMed:[7721776](#)). Lewis-positive (Le(+)) individuals have an active enzyme while Lewis-negative (Le(-)) individuals have an inactive enzyme (PubMed:[1977660](#)).

Cellular Location

Golgi apparatus, Golgi stack membrane; Single- pass type II membrane protein
Note=Membrane-bound form in trans cisternae of Golgi

Tissue Location

Highly expressed in stomach, colon, small intestine, lung and kidney and to a lesser extent in salivary gland, bladder, uterus and liver

BG8,Lewisy - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BG8,Lewisy - Images