

**Anti-LAG3 / CD223 Reference Antibody (relatlimab)  
Recombinant Antibody  
Catalog # APR10029****Specification**

---

**Anti-LAG3 / CD223 Reference Antibody (relatlimab) - Product Information**

Application	FC, Kinetics, Animal Model
Primary Accession	<a href="#">P18627</a>
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4SP
Calculated MW	145.58 KDa

**Anti-LAG3 / CD223 Reference Antibody (relatlimab) - Additional Information****Target/Specificity**  
LAG3 / CD223**Endotoxin**  
< 0.001EU/ µg,determined by LAL method.**Conjugation**  
Unconjugated**Expression system**  
CHO Cell**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-LAG3 / CD223 Reference Antibody (relatlimab) - Protein Information****Name** LAG3 ([HGNC:6476](#))**Synonyms** FDC**Function**  
Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed:<a href="http://www.uniprot.org/citations/20421648" target="\_blank">20421648</a>, PubMed:<a href="http://www.uniprot.org/citations/7805750" target="\_blank">7805750</a>, PubMed:<a href="http://www.uniprot.org/citations/8647185" target="\_blank">8647185</a>). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and

CD4(+) T-cells (PubMed:<a href="http://www.uniprot.org/citations/20421648" target="\_blank">20421648</a>, PubMed:<a href="http://www.uniprot.org/citations/7805750" target="\_blank">7805750</a>, PubMed:<a href="http://www.uniprot.org/citations/8647185" target="\_blank">8647185</a>). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:<a href="http://www.uniprot.org/citations/8647185" target="\_blank">8647185</a>).

#### Cellular Location

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

### Anti-LAG3 / CD223 Reference Antibody (relatlimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-LAG3 / CD223 Reference Antibody (relatlimab) - Images




Anti-LAG3 / CD223 Reference Antibody (relatlimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%




The purity of Anti-LAG3 / CD223 Reference Antibody (relatlimab) is more than 100% ,determined by SEC-HPLC.



Immobilized human LAG3 His at 2 µg/mL can bind Anti-LAG3 / CD223 Reference Antibody (relatlimab)  $EC_{50}=0.02812$  µg/mL.



Human LAG3 CHO cells were stained with Anti-LAG3 / CD223 Reference Antibody (relatlimab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS,  $EC_{78}=0.5464$  µg/mL



Relatlimab inhibited the tumor growth of MC38 on Human LAG3 mice. The result showed significant anti-tumor effects, with an tumor inhibition rate (TGI) of 53.8% at 10 mpk at D28.