

# Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3)

Recombinant Antibody Catalog # APR10043

## **Specification**

# Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) - Product Information

Application FC, Kinetics, Animal Model

Primary Accession
Reactivity
Clonality
Isotype
Calculated MW

P49682
Human
Monoclonal
IgG1
150 KDa

# Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) - Additional Information

Target/Specificity CXCR3 / GPR9 / CD183

**Endotoxin** 

< 0.001EU/ μg, determined by LAL method.

**Conjugation** Unconjugated

**Expression system** 

CHO Cell

#### **Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

# Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) - Protein Information

Name CXCR3

**Synonyms** GPR9

#### **Function**

[Isoform 1]: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11 and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G- protein signaling pathway (PubMed:<a

href="http://www.uniprot.org/citations/12782716" target="\_blank">12782716</a>). Binds to CCL21. Probably promotes cell chemotaxis response. Upon activation by PF4, induces activated T-lymphocytes migration mediated via downstream Ras/extracellular signal-regulated kinase (ERK) signaling. [Isoform 3]: Mediates the activity of CXCL11.



## **Cellular Location**

[Isoform 1]: Cell membrane; Multi-pass membrane protein

#### **Tissue Location**


Isoform 1 and isoform 2 are mainly expressed in heart, kidney, liver and skeletal muscle. Isoform 1 is also expressed in placenta. Isoform 2 is expressed in endothelial cells. Expressed in T-cells (at protein level).

## Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) - Images




Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) is more than 100% ,determined by SEC-HPLC.





Immobilized human CXCR3 VLP Protein at 8  $\mu$ g/mL can bind Anti-CXCR3 / GPR9 / CD183 Reference Antibody (Genzyme patent anti-CXCR3) $\square$ EC50=0.173  $\mu$ g/mL