

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab)

Recombinant Antibody Catalog # APR10265

Specification

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) - Product Information

Application Primary Accession Reactivity Clonality Isotype

Calculated MW

FC, Kinetics, Animal Model

<u>Q9H7M9</u>

Human, Mouse Monoclonal

IqG1

144.52 KDa

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) - Additional Information

Target/Specificity B7-H5 / VISTA

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) - Protein Information

Name VSIR (HGNC:30085)

Function

Immunoregulatory receptor which inhibits the T-cell response (PubMed:24691993). May promote differentiation of embryonic stem cells, by inhibiting BMP4 signaling (By similarity). May stimulate MMP14- mediated MMP2 activation (PubMed:20666777).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in spleen. Detected on a number of myeloid cells including CD11b monocytes, CD66b+ neutrophils, at low levels on CD4+ and CD8+ T-cells, and in a subset of NK cells. Not detected on



Tel: 858.875.1900 Fax: 858.875.1999


B cells (at protein level). Expressed at high levels in placenta, spleen, plasma blood leukocytes, and lung. Expressed at moderate levels in lymph node, bone marrow, fat, uterus, and trachea Has low expression levels in other tissues

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) - Images




Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%




The purity of Anti-B7-H5 / VISTA Reference Antibody (onvatilimab)is more than 98.11% ,determined by SEC-HPLC.





Immobilized human B7-H5 / VISTA His at 2 $\,\mu g/mL$ can bind Anti-B7-H5 / VISTA Reference Antibody (onvatilimab)[]EC50=0.01513 $\mu g/mL$



Human B7H5 CHO-K cells were stained with Anti-B7-H5 / VISTA Reference Antibody (onvatilimab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC328=0.9471 μ g/mL