

**Anti-ICOS / CD278 Reference Antibody (vopratelimab)  
Recombinant Antibody  
Catalog # APR10287****Specification**

---

**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Product Information**

|                   |                          |
|-------------------|--------------------------|
| Application       | FC, E, FTA               |
| Primary Accession | <a href="#">O9Y6W8</a>   |
| Reactivity        | Cynomolgus, Human, Mouse |
| Clonality         | Monoclonal               |
| Isotype           | IgG1                     |
| Calculated MW     | 145.8 KDa                |

**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Additional Information****Target/Specificity**  
ICOS / CD278**Endotoxin**  
< 0.001EU/ µg,determined by LAL method.**Conjugation**  
Unconjugated**Expression system**  
CHO Cell**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Storage**  
-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Protein Information****Name** ICOS**Synonyms** AILIM**Function**  
Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin

isotypes (By similarity).

#### Cellular Location

[Isoform 1]: Cell membrane; Single- pass type I membrane protein

#### Tissue Location


Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

### Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Images



Anti-ICOS / CD278 Reference Antibody (vopratelimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-ICOS / CD278 Reference Antibody (vopratelimab) is more than 99.09%, determined by SEC-HPLC.