

**Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab)
Recombinant Antibody
Catalog # APR10296****Specification****Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) - Product Information**

Application	FC, E, FTA
Primary Accession	P21980
Reactivity	Rat, Cynomolgus, Human
Clonality	Monoclonal
Isotype	IgG4SP
Calculated MW	145 KDa

Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) - Additional Information**Target/Specificity**

TGM2 / Transglutaminase 2

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Storage

-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.

Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) - Protein Information**Name** TGM2 {ECO:0000303|PubMed:17939176, ECO:0000312|HGNC:HGNC:11778}**Function**

Calcium-dependent acyltransferase that catalyzes the formation of covalent bonds between peptide-bound glutamine and various primary amines, such as gamma-amino group of peptide-bound lysine, or mono- and polyamines, thereby producing cross-linked or aminated proteins, respectively (PubMed:9252372, PubMed:23941696, PubMed:31991788). Involved in many biological processes, such as bone

development, angiogenesis, wound healing, cellular differentiation, chromatin modification and apoptosis (PubMed: [1683874](http://www.uniprot.org/citations/1683874), PubMed: [7935379](http://www.uniprot.org/citations/7935379), PubMed: [9252372](http://www.uniprot.org/citations/9252372), PubMed: [27270573](http://www.uniprot.org/citations/27270573)). Acts as a protein-glutamine gamma- glutamyltransferase by mediating the cross-linking of proteins, such as ACO2, HSPB6, FN1, HMGB1, RAP1GDS1, SLC25A4/ANT1, SPP1 and WDR54 (PubMed: [23941696](http://www.uniprot.org/citations/23941696), PubMed: [24349085](http://www.uniprot.org/citations/24349085), PubMed: [29618516](http://www.uniprot.org/citations/29618516), PubMed: [30458214](http://www.uniprot.org/citations/30458214)). Under physiological conditions, the protein cross-linking activity is inhibited by GTP; inhibition is relieved by Ca(2+) in response to various stresses (PubMed: [7649299](http://www.uniprot.org/citations/7649299), PubMed: [7592956](http://www.uniprot.org/citations/7592956), PubMed: [18092889](http://www.uniprot.org/citations/18092889)). When secreted, catalyzes cross-linking of proteins of the extracellular matrix, such as FN1 and SPP1 resulting in the formation of scaffolds (PubMed: [12506096](http://www.uniprot.org/citations/12506096)). Plays a key role during apoptosis, both by (1) promoting the cross-linking of cytoskeletal proteins resulting in condensation of the cytoplasm, and by (2) mediating cross-linking proteins of the extracellular matrix, resulting in the irreversible formation of scaffolds that stabilize the integrity of the dying cells before their clearance by phagocytosis, thereby preventing the leakage of harmful intracellular components (PubMed: [7935379](http://www.uniprot.org/citations/7935379), PubMed: [9252372](http://www.uniprot.org/citations/9252372)). In addition to protein cross-linking, can use different monoamine substrates to catalyze a vast array of protein post-translational modifications: mediates aminylation of serotonin, dopamine, noradrenaline or histamine into glutamine residues of target proteins to generate protein serotonylation, dopaminylation, noradrenalinylation or histaminylation, respectively (PubMed: [23797785](http://www.uniprot.org/citations/23797785), PubMed: [30867594](http://www.uniprot.org/citations/30867594)). Mediates protein serotonylation of small GTPases during activation and aggregation of platelets, leading to constitutive activation of these GTPases (By similarity). Plays a key role in chromatin organization by mediating serotonylation and dopaminylation of histone H3 (PubMed: [30867594](http://www.uniprot.org/citations/30867594), PubMed: [32273471](http://www.uniprot.org/citations/32273471)). Catalyzes serotonylation of 'Gln-5' of histone H3 (H3Q5ser) during serotonergic neuron differentiation, thereby facilitating transcription (PubMed: [30867594](http://www.uniprot.org/citations/30867594)). Acts as a mediator of neurotransmission-independent role of nuclear dopamine in ventral tegmental area (VTA) neurons: catalyzes dopaminylation of 'Gln- 5' of histone H3 (H3Q5dop), thereby regulating relapse-related transcriptional plasticity in the reward system (PubMed: [32273471](http://www.uniprot.org/citations/32273471)). Regulates vein remodeling by mediating serotonylation and subsequent inactivation of ATP2A2/SERCA2 (By similarity). Also acts as a protein deamidase by mediating the side chain deamidation of specific glutamine residues of proteins to glutamate (PubMed: [9623982](http://www.uniprot.org/citations/9623982), PubMed: [20547769](http://www.uniprot.org/citations/20547769)). Catalyzes specific deamidation of protein gliadin, a component of wheat gluten in the diet (PubMed: [9623982](http://www.uniprot.org/citations/9623982)). May also act as an isopeptidase cleaving the previously formed cross-links (PubMed: [26250429](http://www.uniprot.org/citations/26250429), PubMed: [27131890](http://www.uniprot.org/citations/27131890)). Also able to participate in signaling pathways independently of its acyltransferase activity: acts as a signal transducer in alpha-1 adrenergic receptor-mediated stimulation of phospholipase C-delta (PLCD) activity and is required for coupling alpha-1 adrenergic agonists to the stimulation of phosphoinositide lipid metabolism (PubMed: [8943303](http://www.uniprot.org/citations/8943303)

target="_blank">8943303).

Cellular Location


Cytoplasm, cytosol. Nucleus. Chromosome. Secreted, extracellular space, extracellular matrix. Cell membrane {ECO:0000250|UniProtKB:Q9WVJ6}. Mitochondrion. Note=Mainly localizes to the cytosol (PubMed:9575137). Present at much lower level in the nucleus and chromatin (PubMed:9575137). Also secreted via a non-classical secretion pathway to the extracellular matrix (PubMed:27270573)

Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) - Images



Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-TGM2 / Transglutaminase 2 Reference Antibody (zampilimab) is more than 95%, determined by SEC-HPLC.