

**Anti-IL-5 Reference Antibody (depemokimab)
Recombinant Antibody
Catalog # APR10546****Specification**

Anti-IL-5 Reference Antibody (depemokimab) - Product Information

Application	FC, E, FTA
Primary Accession	P05113
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.8 KDa

Anti-IL-5 Reference Antibody (depemokimab) - Additional Information**Target/Specificity**

IL-5

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Storage

-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.

Anti-IL-5 Reference Antibody (depemokimab) - Protein Information**Name** IL5**Function**

Homodimeric cytokine expressed predominantly by T-lymphocytes and NK cells that plays an important role in the survival, differentiation, and chemotaxis of eosinophils (PubMed:2653458, PubMed:9010276). Acts also on activated and resting B-cells to induce immunoglobulin production, growth, and differentiation (By similarity). Mechanistically, exerts its biological effects through a receptor composed of IL5RA subunit and the cytokine receptor common subunit beta/CSF2RB (PubMed:1495999, PubMed:1495999).

href="http://www.uniprot.org/citations/22528658" target="_blank">22528658). Binding to the receptor leads to activation of various kinases including LYN, SYK and JAK2 and thereby propagates signals through the RAS-MAPK and JAK-STAT5 pathways respectively (PubMed:7613138).

Cellular Location


Secreted.

Anti-IL-5 Reference Antibody (depemokimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-5 Reference Antibody (depemokimab) - Images



Anti-IL-5 Reference Antibody (depemokimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IL-5 Reference Antibody (depemokimab) is more than 98.75%, determined by SEC-HPLC.