

**Anti-ALCAM / CD166 Reference Antibody (AT002)
Recombinant Antibody
Catalog # APR10622**

Specification

Anti-ALCAM / CD166 Reference Antibody (AT002) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	Q13740
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	150 KDa

Anti-ALCAM / CD166 Reference Antibody (AT002) - Additional Information

Target/Specificity

ALCAM / CD166

Endotoxin

< 0.001EU/ µg, determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-ALCAM / CD166 Reference Antibody (AT002) - Protein Information

Name ALCAM

Synonyms MEMD {ECO:0000303|PubMed:9502422}

Function

Cell adhesion molecule that mediates both heterotypic cell-cell contacts via its interaction with CD6, as well as homotypic cell-cell contacts (PubMed:15048703, PubMed:15496415, PubMed:16352806, PubMed:23169771, PubMed:24945728, PubMed:7760007). Promotes T-cell activation and proliferation via its interactions with CD6 (PubMed:15048703, PubMed:>16352806, PubMed:>24945728). Contributes to the formation and maturation of the immunological synapse via its interactions with CD6 (PubMed:>15294938, PubMed:>16352806). Mediates homotypic interactions with cells that express ALCAM (PubMed:>15496415, PubMed:>16352806). Acts as a ligand for the LILRB4 receptor, enhancing LILRB4-mediated inhibition of T cell proliferation (PubMed:>29263213). Required for normal hematopoietic stem cell engraftment in the bone marrow (PubMed:>24740813). Mediates attachment of dendritic cells onto endothelial cells via homotypic interaction (PubMed:>23169771). Inhibits endothelial cell migration and promotes endothelial tube formation via homotypic interactions (PubMed:>15496415, PubMed:>23169771). Required for normal organization of the lymph vessel network. Required for normal hematopoietic stem cell engraftment in the bone marrow. Plays a role in hematopoiesis; required for normal numbers of hematopoietic stem cells in bone marrow. Promotes in vitro osteoblast proliferation and differentiation (By similarity). Promotes neurite extension, axon growth and axon guidance; axons grow preferentially on surfaces that contain ALCAM. Mediates outgrowth and pathfinding for retinal ganglion cell axons (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:Q61490}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q61490}. Note=Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with CD6 and the TCR/CD3 complex at the immunological synapse (PubMed:15294938).

Tissue Location


Detected on hematopoietic stem cells derived from umbilical cord blood (PubMed:24740813). Detected on lymph vessel endothelial cells, skin and tonsil (PubMed:23169771). Detected on peripheral blood monocytes (PubMed:15048703). Detected on monocyte- derived dendritic cells (at protein level) (PubMed:16352806). Detected at low levels in spleen, placenta, liver (PubMed:9502422). Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells (PubMed:7760007). Isoform 1 and isoform 3 are detected in vein and artery endothelial cells, astrocytes, keratinocytes and artery smooth muscle cells (PubMed:15496415). Expressed by neurons in the brain Restricted expression in tumor cell lines. Detected in highly metastasizing melanoma cell lines (PubMed:9502422)

Anti-ALCAM / CD166 Reference Antibody (AT002) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ALCAM / CD166 Reference Antibody (AT002) - Images



Anti-ALCAM / CD166 Reference Antibody (AT002) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-ALCAM / CD166 Reference Antibody (AT002) is more than 95% ,determined by SEC-HPLC.