

**Anti-LINGO1 Reference Antibody (opicinumab)
Recombinant Antibody
Catalog # APR10684****Specification**

Anti-LINGO1 Reference Antibody (opicinumab) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	Q96FE5
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	144.76 KDa

Anti-LINGO1 Reference Antibody (opicinumab) - Additional Information**Target/Specificity**
LINGO1**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-LINGO1 Reference Antibody (opicinumab) - Protein Information****Name** LINGO1**Synonyms** LERN1, LRRN6A**Function**
Functional component of the Nogo receptor signaling complex (RTN4R/NGFR) in RhoA activation responsible for some inhibition of axonal regeneration by myelin-associated factors (PubMed:14966521, PubMed:15694321). Is also an important negative regulator of oligodendrocyte differentiation and axonal myelination (PubMed:15895088). Acts in conjunction with RTN4 and RTN4R in regulating neuronal precursor cell motility during cortical development (By similarity).**Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:Q9D1T0}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9D1T0}

Tissue Location


Expressed exclusively in the central nervous system. Highest level in the in amygdala, hippocampus, thalamus and cerebral cortex. In the rest of the brain a basal expression seems to be always present. Up-regulated in substantia nigra neurons from Parkinson disease patients.

Anti-LINGO1 Reference Antibody (opicinumab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-LINGO1 Reference Antibody (opicinumab) - Images



Anti-LINGO1 Reference Antibody (opicinumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-LINGO1 Reference Antibody (opicinumab) is more than 99.03%, determined by SEC-HPLC.