

# Anti-DKK1 Reference Antibody (BHQ880)

Recombinant Antibody Catalog # APR10881

### Specification

## Anti-DKK1 Reference Antibody (BHQ880) - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW FC, Kinetics, Animal Model <u>094907</u> Human Monoclonal IgG1 142.96 KDa

### Anti-DKK1 Reference Antibody (BHQ880) - Additional Information

Target/Specificity DKK1

**Endotoxin** < 0.001EU/ μg,determined by LAL method.

Conjugation Unconjugated

Expression system CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

### Anti-DKK1 Reference Antibody (BHQ880) - Protein Information

Name DKK1

#### Function

Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6 (PubMed:<a href="http://www.uniprot.org/citations/22000856" target="\_blank">22000856</a>). DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (PubMed:<a href="http://www.uniprot.org/citations/17143291" target="\_blank">17143291</a>). Inhibits the pro-apoptotic function of KREMEN1 in a Wnt-independent manner, and has anti-apoptotic activity (By similarity).

Cellular Location Secreted.




Tissue Location Placenta.

#### Anti-DKK1 Reference Antibody (BHQ880) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### Anti-DKK1 Reference Antibody (BHQ880) - Images



Anti-DKK1 Reference Antibody (BHQ880) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-DKK1 Reference Antibody (BHQ880)is more than 100% ,determined by SEC-HPLC.