

**Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1)
Recombinant Antibody
Catalog # APR10926****Specification**

Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	O60565
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4SP
Calculated MW	150 KDa

Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) - Additional Information**Target/Specificity**

GREM1 / Gremlin

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) - Protein Information**Name** GREM1**Synonyms** CKTSF1B1, DAND2, DRM**Function**

Cytokine that may play an important role during carcinogenesis and metanephric kidney organogenesis, as a BMP antagonist required for early limb outgrowth and patterning in maintaining the FGF4-SHH feedback loop. Down-regulates the BMP4 signaling in a dose-dependent manner (By similarity). Antagonist of BMP2; inhibits BMP2-mediated differentiation of osteoblasts (in vitro) (PubMed:27036124). Acts as inhibitor of monocyte chemotaxis. Can inhibit the growth or viability of normal cells but not transformed cells when is overexpressed (By similarity).


Cellular Location
Secreted.**Tissue Location**

Highly expressed in small intestine, fetal brain and colon. Expression is restricted to intestinal subepithelial myofibroblasts (ISEMFs) at the crypt base. In subjects with HGPS, by contrast, GREM1 is expressed, not only in basal ISEMFs, but also at very high levels in epithelial cells (predominantly colonocytes), with expression extending most of the way up the sides of the crypt. Weakly expressed in brain, ovary, prostate, pancreas and skeletal muscle. In brain found in the region localized around the internal capsule in the large subcortical nuclei, including caudate, putamen, substantia nigra, thalamus and subthalamus. Predominantly expressed in normal cells including neurons, astrocytes and fibroblasts


Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) - Images

Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-GREM1 / Gremlin Reference Antibody (Regeneron patent anti-GREM1) is more than 95% ,determined by SEC-HPLC.