

Anti-IGF-2 Reference Antibody (DX-2647)

Recombinant Antibody Catalog # APR10947

Specification

Anti-IGF-2 Reference Antibody (DX-2647) - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW FC, Kinetics, Animal Model <u>P01344</u> Human Monoclonal IgG1 145.78 KDa

Anti-IGF-2 Reference Antibody (DX-2647) - Additional Information

Target/Specificity IGF-2

Endotoxin < 0.001EU/ μg,determined by LAL method.

Conjugation Unconjugated

Expression system CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-IGF-2 Reference Antibody (DX-2647) - Protein Information

Name IGF2 (HGNC:5466)

Function

The insulin-like growth factors possess growth-promoting activity (By similarity). Major fetal growth hormone in mammals. Plays a key role in regulating fetoplacental development. IGF2 is influenced by placental lactogen. Also involved in tissue differentiation. In adults, involved in glucose metabolism in adipose tissue, skeletal muscle and liver (Probable). Acts as a ligand for integrin which is required for IGF2 signaling (PubMed:28873464). Positively regulates myogenic transcription factor MYOD1 function by facilitating the recruitment of transcriptional coactivators, thereby controlling muscle terminal differentiation (By similarity). Inhibits myoblast differentiation and modulates metabolism via increasing the mitochondrial respiration rate (By similarity).

Cellular Location Secreted.



Tissue Location

Expressed in heart, placenta, lung, liver, muscle, kidney, tongue, limb, eye and pancreas.

Anti-IGF-2 Reference Antibody (DX-2647) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

Anti-IGF-2 Reference Antibody (DX-2647) - Images

kDa	NR	М	R
			1
185			
115			
80		-	
65		-	
50		-	
30		-	-
25			
15		Ξ.	1
10	-		

Anti-IGF-2 Reference Antibody (DX-2647) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IGF-2 Reference Antibody (DX-2647)is more than 95% ,determined by SEC-HPLC.