

**Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1)
Recombinant Antibody
Catalog # APR10953****Specification**

Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	Q9UHF4
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Calculated MW	150 KDa

Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) - Additional Information**Target/Specificity**

IL-20Ra

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) - Protein Information**Name** IL20RA**Function**

The IL20RA/IL20RB dimer is a receptor for IL19, IL20 and IL24. The IL20RA/IL10RB dimer is a receptor for IL26.

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location


Widely expressed with highest levels in skin and testis and high levels in brain. Highly expressed in psoriatic skin

Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) - Images



Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-IL-20Ra Reference Antibody (Cheng Kung U. patent anti-IL-20R1) is more than 95%, determined by SEC-HPLC.